

Before

Existing Label

Allergy Tablets

INDICATIONS: Provides effective, temporary relief of sneezing, watery and itchy eyes, and runny nose due to hay fever and other upper respiratory allergies.

DIRECTIONS: Adults and children 12 years and over—1 tablet every 4 to 6 hours, not to exceed 6 tablets in 24 hours or as directed by a physician. Children 6 to 11 years—one half the adult dose (break tablet in half) every 4 to 6 hours, not to exceed 3 whole tablets in 24 hours. For children under 6 years, consult a physician.

EACH TABLET CONTAINS: Chlorpheniramine Maleate 4 mg. **May also contain** (may differ from brand): D&C Yellow No. 10, Lactose, Magnesium Stearate, Microcrystalline Cellulose, Pregelatinized Starch.


WARNINGS: May cause excitability especially in children. Do not take this product unless directed by a physician, if you have a breathing problem such as emphysema or chronic bronchitis, or if you have glaucoma or difficulty in urination due to enlargement of the prostate gland. May cause drowsiness; alcohol, sedatives and tranquilizers may increase the drowsiness effect. Avoid alcoholic beverages, and do not take this product if you are taking sedatives or tranquilizers without first consulting your physician. Use caution when driving a motor vehicle or operating machinery. As with any drug, if you are pregnant or nursing a baby, seek the advice of a health professional before using this product. Keep this and all drugs out of the reach of children. In case of accidental overdose, seek professional assistance or contact a Poison Control Center immediately.

Store at controlled room temperature 2°-30°C (36°-86°F).

Use by expiration date printed on package.

Protect from excessive moisture.

For better identification keep tablets in carton until used.



N
3 111111 11111 8

Made in U.S.A.

After

New Standard Labeling Format

| Drug Facts | |
|---|--|
| Active ingredient (in each tablet) | Purpose |
| Chlorpheniramine maleate 2 mg..... | Antihistamine |
| Uses temporarily relieves these symptoms due to hay fever or other upper respiratory allergies: ■ sneezing ■ runny nose ■ itchy, watery eyes ■ itchy throat | |
| Warnings Ask a doctor before use if you have ■ glaucoma ■ a breathing problem such as emphysema or chronic bronchitis ■ trouble urinating due to an enlarged prostate gland Ask a doctor or pharmacist before use if you are taking tranquilizers or sedatives | |
| When using this product ■ drowsiness may occur ■ avoid alcoholic drinks ■ alcohol, sedatives, and tranquilizers may increase drowsiness ■ be careful when driving a motor vehicle or operating machinery ■ excitability may occur, especially in children | |
| If pregnant or breast-feeding, ask a health professional before use. Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away. | |
| Directions | |
| adults and children 12 years and over | take 2 tablets every 4 to 6 hours; not more than 12 tablets in 24 hours |
| children 6 years to under 12 years | take 1 tablet every 4 to 6 hours; not more than 6 tablets in 24 hours |
| children under 6 years | ask a doctor |

| | |
|---|--|
| Drug Facts (continued) | |
| Other information ■ store at 20-25°C (68-77°F) ■ protect from excessive moisture | |
| Inactive ingredients D&C yellow no. 10, lactose, magnesium stearate, microcrystalline cellulose, pregelatinized starch | |