

Medication Guide

Medication Guide ILARIS (i-LAHR-us) (canakinumab) injection for subcutaneous use

What is the most important information I should know about ILARIS?

ILARIS can cause serious side effects, including:

- **Increased risk of serious infections.** ILARIS can lower the ability of your immune system to fight infections. Your healthcare provider should:
 - test you for tuberculosis (TB) before you receive ILARIS
 - monitor you closely for symptoms of TB during treatment with ILARIS
 - check you for symptoms of any type of infection before, during, and after your treatment with ILARIS

Tell your healthcare provider right away if you have any symptoms of an infection such as fever, sweats or chills, cough, flu-like symptoms, weight loss, shortness of breath, blood in your phlegm, sores on your body, warm or painful areas on your body, diarrhea or stomach pain, or feeling very tired.

See "What are possible side effects of ILARIS?" for more information about side effects.

What is ILARIS?

ILARIS is a prescription medicine injected by your healthcare provider just below the skin (subcutaneous) used to treat:

- Adults and children 4 years of age and older who have auto-inflammatory diseases called Cryopyrin-Associated Periodic Syndromes (CAPS), including:
 - Familial Cold Auto-inflammatory Syndrome (FCAS)
 - Muckle-Wells Syndrome (MWS)
- Systemic Juvenile Idiopathic Arthritis (SJIA) in children 2 years of age and older.

It is not known if ILARIS is safe and effective when used to treat SJIA in children under 2 years of age or when used to treat CAPS in children under 4 years of age.

Who should not receive ILARIS?

- Do not receive ILARIS if you are allergic to canakinumab or any of the ingredients in ILARIS. See the end of this Medication Guide for a complete list of ingredients in ILARIS.

What should I tell my healthcare provider before receiving ILARIS?

Before you receive ILARIS, tell your healthcare provider about all your medical conditions, including if you:

- think you have or are being treated for an active infection
- have symptoms of an infection
- have a history of infections that keep coming back
- have a history of low white blood cells
- have or have had HIV, Hepatitis B, or Hepatitis C
- are scheduled to receive any immunizations (vaccines). You should not get 'live vaccines' if you are receiving ILARIS.
- are pregnant or planning to become pregnant. It is not known if ILARIS will harm your unborn baby. Tell your healthcare provider right away if you become pregnant while receiving ILARIS.
- are breastfeeding or planning to breastfeed. It is not known if ILARIS passes into your breast milk. You and your healthcare provider should decide if you will receive ILARIS or breastfeed. You should not do both.

Tell your healthcare provider about all the medicines you take, including prescription and nonprescription medicines, vitamins, and herbal supplements. Especially tell your healthcare provider if you take:

- Medicines that affect your immune system
- Medicines called IL-1 blocking agents such as Kineret[®] (anakinra), Arcalyst[®] (rilonacept)
- Medicines called Tumor Necrosis Factor (TNF) inhibitors such as Enbrel[®] (etanercept), Humira[®] (adalimumab), Remicade[®] (infliximab), Simponi[®] (golimumab), or Cimzia[®] (certolizumab pegol)
- Medicines that effect enzyme metabolism

Ask your healthcare provider if you are not sure.

How should I receive ILARIS?

- ILARIS is given by your healthcare provider every 8 weeks for CAPS and every 4 weeks for SJIA.

What are the possible side effects of ILARIS?

ILARIS can cause serious side effects, including:

- See **“What is the most important information I should know about ILARIS?”**
- **decreased ability of your body to fight infections (immunosuppression).** For

people treated with medicines that cause immunosuppression like ILARIS, the chances of getting cancer may increase.

- **allergic reactions.** Allergic reactions can happen while you are receiving ILARIS. Call your healthcare provider right away if you have any of these symptoms of an allergic reaction:
 - rash
 - itching and hives
 - difficulty breathing or swallowing
 - dizziness or feeling faint
- **risk of infection with live vaccines.** You should not get live vaccines if you are receiving ILARIS. Tell your healthcare provider if you are scheduled to receive any vaccines.

The most common side effects of ILARIS include:

When ILARIS is used for the treatment of CAPS:

- cold symptoms
- diarrhea
- flu (influenza)
- runny nose
- headache
- cough
- body aches
- feeling like you are spinning (vertigo)
- weight gain
- injection site reactions (such as redness, swelling, warmth, or itching)
- nausea

When ILARIS is used for treatment of SJIA:

- cold symptoms
- upper respiratory tract infection
- pneumonia
- runny nose
- sore throat
- urinary tract infection
- nausea, vomiting, and diarrhea (gastroenteritis)
- stomach pain
- injection site reactions

Tell your healthcare provider about any side effect that bothers you or does not go away. These are not all the possible side effects of ILARIS. For more information, ask your healthcare provider or pharmacist.

Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088.

General information about the safe and effective use of ILARIS.

Medicines are sometimes prescribed for purposes other than those listed in this Medication Guide. Do not use ILARIS for a condition for which it was not prescribed. Do not give ILARIS to other people, even if they have the same condition as you. It may harm them.

This Medication Guide summarizes the most important information about ILARIS. If you would like more information, talk with your healthcare provider. You can ask your healthcare provider or pharmacist for information about ILARIS that was written for health professionals. For more information about ILARIS, call 1-877-452-7471 or visit www.ILARIS.com.

What are the ingredients in ILARIS?

Active ingredient: canakinumab

Inactive ingredients: sucrose, L-histidine, L-histidine HCl monohydrate, polysorbate 80, preservative-free sterile water for injection

This Medication Guide has been approved by the U.S. Food and Drug Administration.

Distributed by:
Novartis Pharmaceuticals Corporation
East Hanover, New Jersey 07936

© Novartis

Kineret[®], Arcalyst[®], Enbrel[®], Humira[®], Remicade[®], Simponi[®], and Cimzia[®] are trademarks of Amgen, Regeneron, Immunex Corporation, AbbVie Biotechnology Ltd., Centocor Ortho Biotech Inc., Janssen Biotech Inc., and the UCB Group of companies, respectively.

T2016-07/T2016-07
July 2016/ July 2016