

NICARBAZIN/BACITRACIN METHYLENE DISALICYLATE  
TYPE C

BLUE BIRD  
CHICKEN FEED  
MEDICATED

**INDICATIONS FOR USE**

Broiler chickens; aid in preventing outbreaks of cecal (*Eimeria tenella*) and intestinal (*E. acervulina*, *E. maxima*, *E. necatrix*, *E. brunetti*) coccidiosis; for increased rate of weight gain and improved feed efficiency.

**ACTIVE DRUG INGREDIENTS**

Bacitracin methylene disalicylate..... 30g/ton  
Nicarbazin ..... 90.8 -181.6g/ton (0.01 – 0.02%)\*

**GUARANTEED ANALYSIS**

Crude Protein (Min) ..... %  
Lysine (Min)..... %  
Methionine (Min) ..... %  
Crude Fat (Min)..... %  
Crude Fiber (Max)..... %  
Calcium (Min) ..... %  
Calcium (Max) ..... %  
Phosphorus (Min) ..... %  
Salt<sup>a</sup> ..... %  
Sodium<sup>b</sup> ..... %

<sup>a</sup> Guarantee required only when nutrient added except when the feed is intended, represented or serves as the principal source of the nutrient.

<sup>b</sup> Sodium guarantee required only when total sodium exceeds that furnished by the maximum salt guarantee.

**INGREDIENTS**

Ingredients as defined by AAFCO.

**FEEDING DIRECTIONS**

Feed continuously as the sole ration from the time chicks are placed on litter until past the time when coccidiosis is ordinarily a hazard.

**WARNING:** Withdraw 4 days before slaughter for use levels at or below 113.5 g nicarbazin/ton ( $\leq 0.0125\%$ ).  
Withdraw 5 days before slaughter for use levels above 113.5 g nicarbazin/ton ( $> 0.0125\%$ ).

**CAUTION:** Do not use as a treatment for coccidiosis. Do not feed to laying hens. Do not use in flushing mashes. If losses exceed 0.5 percent in a 2-day period, obtain an accurate diagnosis and follow the instructions of your veterinarian or poultry pathologist. Broilers fed feed medicated with nicarbazin may show reduced heat tolerance when exposed to high temperature and high humidity to which they have not been accustomed and under severe conditions, fatalities may result. An ample supply of drinking water and adequate ventilation will improve the birds' tolerance of heat.

\_\_\_ lb (\_\_\_ kg) NET WEIGHT

**BLUEBIRD FEED MILL**  
Anytown, USA 12345

\* The finished feed should contain a specific concentration of nicarbazin, and the WARNING withdrawal statement should reflect this use level.