

NICARBAZIN
TYPE C

BLUE BIRD
CHICKEN FEED
MEDICATED

Aid in preventing outbreaks of cecal (*Eimeria tenella*) and intestinal (*E. acervulina*, *E. maxima*, *E. necatrix*, and *E. brunetti*) coccidiosis.

ACTIVE DRUG INGREDIENT

Nicarbazin90.8 - 181.6 g/ton^a

GUARANTEED ANALYSIS

Crude Protein (Min)..... _____ %
Lysine (Min)..... _____ %
Methionine (Min) _____ %
Crude Fat (Min)..... _____ %
Crude Fiber (Min)..... _____ %
Calcium (Min) _____ %
Calcium (Max)..... _____ %
Phosphorus (Min) _____ %
Salt, if added (Min)..... _____ %
Salt, if added (Max)..... _____ %
Total Sodium (Min)*..... _____ %
Total Sodium (Max)* _____ %

* Minimum and maximum percentage of total sodium shall be guaranteed only when total sodium exceeds that furnished by the maximum salt guarantee.

INGREDIENTS

Ingredients as defined by AAFCO.

FEEDING DIRECTIONS

Feed continuously as sole ration from time chicks are placed on litter until past the time when coccidiosis is ordinarily a hazard.

WARNING: Withdraw 4 days before slaughter for use levels at or below 113.5 g nicarbazin/ton.
Withdraw 5 days before slaughter for use levels above 113.5 g nicarbazin/ton.

CAUTION: Do not use as a treatment for coccidiosis. Do not feed to laying hens. Do not use in flushing mashers. If losses exceed 0.5 percent in a 2-day period, obtain an accurate diagnosis and follow the instructions of your veterinarian or poultry pathologist. Broilers fed feed medicated with Nicarb 25% may show reduced heat tolerance when exposed to high temperature and high humidity to which they have not been accustomed and under severe conditions, fatalities may result. An ample supply of drinking water and adequate ventilation will improve the birds' tolerance of heat.

____lb (____kg) NET WEIGHT

BLUE BIRD FEED MILL
Robin, IN 00000

^a The finished feed should contain a specific concentration of nicarbazin, and the WARNING withdrawal statement should reflect this use level.