

**BLUE BIRD  
RBZ  
Type B Medicated Broiler Chicken Feed**

As an aid in the prevention of coccidiosis caused by *Eimeria mivati*, *E. brunetti*, *E. tenella*, *E. acervulina*, *E. maxima*, and *E. necatrix* in broiler chickens only.

**Active Drug Ingredient**

Robenidine hydrochloride.....31 – 3,000 g/ton\*

**Guaranteed Analysis**

Crude Protein (Min) .....	_____	%
Lysine (Min).....	_____	%
Methionine (Min) .....	_____	%
Crude Fat (Min) .....	_____	%
Crude Fiber (Max) .....	_____	%
Calcium (Min).....	_____	%
Calcium (Max).....	_____	%
Phosphorus (Min) .....	_____	%
Salt <sup>a</sup> (Min).....	_____	%
Salt <sup>a</sup> (Max).....	_____	%
Sodium <sup>b</sup> (Min).....	_____	%
Sodium <sup>b</sup> (Max).....	_____	%

<sup>a</sup> Guarantee required only when nutrient added except when the feed is intended, represented or serves as a principal source of the nutrient.

<sup>b</sup> Sodium guarantee required only when total sodium exceeds that furnished by the maximum salt guarantee.

**Ingredients**

Ingredients as defined by AAFCO.

**Mixing Directions**

Thoroughly mix this Type B medicated feed with non-medicated feed ingredients to manufacture one ton of Type C medicated feed for broiler chickens containing 30 grams robenidine per ton. Do not use in feeds containing bentonite.

The following table provides examples of mixing rates:

Type B robenidine concentration (g/ton)	Type B per ton of Type C (lb)	Non-medicated feed per ton of Type C (lb)	Type C robenidine concentration (g/ton)
1,000	60	1,940	30
2,000	30	1,970	30
3,000	20	1,980	30

The resulting Type C medicated feed should be fed continuously as the sole ration. The Type C feed must be fed within 50 days from the date of manufacture.

**Warning**

Do not feed to chickens producing eggs for food. Withdraw 5 days prior to slaughter.

**Manufactured by**  
Blue Bird Feed Mill  
City, State, Zip

**NET WEIGHT ON BAG OR BULK**

\*The final printed feed label should list only a single drug concentration.