

# Blue Bird

Tiamulin Type C Medicated Swine Feed for Control of Swine Dysentery  
STARTER-GROWER-FINISHER FEEDS

**INDICATIONS:**

For control of swine dysentery associated with *Brachyspira* (formerly *Serpulina* or *Treponema*) *hyodysenteriae* susceptible to tiamulin.

**ACTIVE INGREDIENT:**

Tiamulin hydrogen fumarate.....35g/ton

---

**GUARANTEED ANALYSIS**

Crude protein, not less than \_\_\_\_%  
Lysine, not less than \_\_\_\_%  
Crude fat, not less than \_\_\_\_%  
Crude fiber, not more than \_\_\_\_%  
Calcium Min. \_\_\_\_% Max. \_\_\_\_%  
Phosphorus, not less than \_\_\_\_%  
Salt (if added) Min. \_\_\_\_% Max. \_\_\_\_%  
Sodium (if exceeding salt guarantee) \_\_\_\_%  
Selenium \_\_\_\_ppm  
Zinc \_\_\_\_ppm

---

**INGREDIENTS:**

Each ingredient must be specifically named in accordance with the names and definitions adopted by AAFCO.

---

**FEEDING DIRECTIONS:**

Feed continuously as sole ration on premises with a history of swine dysentery but where signs of disease have not yet occurred or following approved treatment of disease. For Use In Animals Only.

**WARNING:**

Withdraw medicated feed 2 days before slaughter following use at 35 g/ton.

**CAUTION:**

1. Use as the only source of tiamulin.
2. For use in swine only. The effects of tiamulin on swine reproductive performance, pregnancy and lactation have not been determined.
3. Swine being treated with tiamulin should not have access to feeds containing polyether ionophores (e.g., lasalocid, monensin, narasin, salinomycin and semduramicin) as adverse reactions may occur.
4. If signs of toxicity occur, discontinue use of medicated feed.

**Manufactured By:**

**Blue Bird Feed Company**  
Robin, IN 99999  
NAH/DEN10-MP/BBC-35/BGL-SD/2  
09/12

**Net Weight:**  
**Lot Number:**