

Blue Bird

Tiamulin Type C Medicated Swine Feed for Treatment of Swine Dysentery
STARTER-GROWER-FINISHER FEEDS

INDICATIONS:

For treatment of swine dysentery associated with *Brachyspira* (formerly *Serpulina* or *Treponema*) *hyodysenteriae* susceptible to tiamulin.

ACTIVE INGREDIENT:

Tiamulin hydrogen fumarate.....200g/ton

GUARANTEED ANALYSIS

Crude protein, not less than ____%
Lysine, not less than ____%
Crude fat, not less than ____%
Crude fiber, not more than ____%
Calcium Min.____% Max.____%
Phosphorus, not less than ____%
Salt (if added) Min.____% Max.____%
Sodium (if exceeding salt guarantee) ____%
Selenium ____ppm
Zinc ____ppm

INGREDIENTS:

Each ingredient must be specifically named in accordance with the names and definitions adopted by AAFCO.

FEEDING DIRECTIONS:

Feed continuously as sole ration for 14 consecutive days. For Use In Animals Only.

WARNING:

Withdraw medicated feed 7 days before slaughter following use at 200 g/ton.

CAUTION:

1. Use as the only source of tiamulin.
2. For use in swine only. The effects of tiamulin on swine reproductive performance, pregnancy and lactation have not been determined.
3. Swine being treated with tiamulin should not have access to feeds containing polyether ionophores (e.g., lasalocid, monensin, narasin, salinomycin and semduramicin) as adverse reactions may occur.
4. If signs of toxicity occur, discontinue use of medicated feed.

Manufactured By:

Blue Bird Feed Company
Robin, IN 99999
NAH/DEN10-MP/BBC-200/BGL-SD/2
09/12

Net Weight:
Lot Number: