

Chlortetracycline Turkey Feed – AUREO^a - SYN
Type B Medicated Feed
(chlortetracycline Type B medicated feed)

Caution: Federal law restricts medicated feed containing this veterinary feed directive (VFD) drug to use by or on the order of a licensed veterinarian.

For control of infectious synovitis caused by *Mycoplasma synoviae* susceptible to chlortetracycline in turkeys.

Active Drug Ingredient

Chlortetracycline^a201 – 80,000 g/ton*

Guaranteed Analysis

Crude Protein (Min)	%
Lysine (Min)	%
Methionine (Min)	%
Crude Fat (Min)	%
Crude Fiber (Max)	%
Calcium (Min)	%
Calcium (Max)	%
Phosphorus (Min)	%
Salt ¹ (Min)	%
Salt ¹ (Max)	%
Sodium ² (Min)	%
Sodium ² (Max)	%

¹ Guarantee required only when nutrient added except when the feed is intended, represented or serves as a principal source of the nutrient.

² Sodium guarantee required only when total sodium exceeds that furnished by the maximum salt guarantee.

Ingredients

Ingredients as defined by AAFCO.

Mixing Directions

Thoroughly mix this Type B medicated feed with non-medicated feed ingredients to manufacture one ton of Type C medicated feed for turkeys containing 200 grams chlortetracycline per ton.

The following table provides examples of mixing rates:

Type B CTC concentration (g/ton)	Type B per ton of Type C (lb)	Non-medicated feed per ton of Type C (lb)	Type C CTC concentration (g/ton)
10,000 (5 g/lb)	40	1960	200
20,000 (10 g/lb)	20	1980	200
40,000 (20 g/lb)	10	1990	200
80,000 (40 g/lb)	5	1995	200

The resulting Type C medicated feed should be fed continuously as the sole ration for 7 to 14 days.

Warning

Do not feed to turkeys producing eggs for human consumption.

Lot Number (if applicable): _____

MANUFACTURED BY

Blue Bird Feed Company
Robin, Indiana 11111

NET WEIGHT ON BAG OR BULK

*The final printed medicate feed label must state a single drug concentration.

^aAureomycin® is the proprietary name of chlortetracycline Type A medicated article (NADA 048-761).

Zoetis Inc. [15 March 2016]