

Ractopamine Medicated Finishing Cattle Feed

Type B Medicated Feed

Do Not Feed Undiluted

IMPORTANT: MUST BE THOROUGHLY MIXED INTO FEED BEFORE USE

For increased rate of weight gain and improved feed efficiency, and for increased rate of weight gain, improved feed efficiency, and increased carcass leanness in cattle fed in confinement for slaughter during the last 28 to 42 days on feed.

Indications	Appropriate Concentration of Ractopamine in Type C Medicated Feed (90% Dry Matter)
Increased Rate of Weight Gain and Improved Feed Efficiency	8.2 to 24.6 g/ton (9 ppm to 27 ppm)
Increased Rate of Weight Gain, Improved Feed Efficiency and Increased Carcass Leanness	9.8 to 24.6 g/ton (11 ppm to 27 ppm)

ACTIVE DRUG INGREDIENT

Ractopamine hydrochloride 26 to 4920 g/ton*
[Up to 4920 g/ton (2.46 g/lb)]

GUARANTEED ANALYSIS

Crude Protein, not less than %
Non-Protein Nitrogen (NPN)¹, not more than..... %
Crude Fat, not less than..... %
Crude Fiber, not more than %
Calcium, not less than %
Calcium, not more than..... %
Phosphorus, not less than..... %
Salt², not less than %
Salt², not more than..... %
Sodium³, not less than..... %
Sodium³, not more than..... %
Potassium, not less than %
Vitamin A^{2,4}, not less than..... I.U./lb

¹When added.

²If added.

³Shall be guaranteed only when total sodium exceeds that furnished by the maximum salt guarantee.

⁴Other than precursors of Vitamin A.

INGREDIENTS

Each ingredient must be named in accordance with the names and definitions adopted by the Association of American Feed Control Officials.

MIXING DIRECTIONS

Thoroughly mix ractopamine Type B Medicated Feed into one ton of complete cattle feed according to the table below to obtain the proper concentration in the Type C Medicated Feed. Prepare an intermediate pre-blend of the medicated feed prior to mixing in a complete feed. Thoroughly mix the required amount in a convenient quantity of feed ingredients, then add to the remaining feed ingredients to make a ton of complete feed. [Use only the portion of the table below on your Type B Medicated Feed product label that is applicable to the concentration of ractopamine in the Type B Medicated Feed which you manufacture.]

Concentration of Ractopamine in Type B Medicated Feed grams/pound	Pounds Type B Medicated Feed To Add Per Ton of Type C Medicated Feed	Resulting Ractopamine Concentration in Type C Medicated Feed ^a grams/ton
0.25	32.8	8.2
	65.6	16.4
	98.4	24.6
0.82	10.0	8.2
	20.0	16.4
	30.0	24.6
1.64	5.0	8.2
	10.0	16.4
	15.0	24.6
2.46	3.3	8.2
	6.6	16.4
	10.0	24.6

^a Based on 90% Dry Matter Basis

WARNING

The active ingredient, ractopamine hydrochloride, is a beta-adrenergic agonist. Individuals with cardiovascular disease should exercise special caution to avoid exposure. Not for use in humans. Keep out of the reach of children. The ractopamine hydrochloride formulation (Type A Medicated Article) poses a low dust potential under usual conditions of handling and mixing. When mixing and handling ractopamine hydrochloride, use protective clothing, impervious gloves, protective eye wear, and a NIOSH-approved dust mask. Operators should wash thoroughly with soap and water after handling. If accidental eye contact occurs, immediately rinse eyes thoroughly with water. If irritation persists, seek medical attention. The material safety data sheet contains more detailed occupational safety information. To report adverse effects, access medical information, or obtain additional product information, call 1-888-963-8471.

CAUTION

Not for animals intended for breeding.

MANUFACTURED BY
BLUE BIRD FEED MILL
Any town, USA 12345

Net Weight lb (kg) on bag or bulk

* Final printed label on formulated Type B medicated feed must bear a single concentration of this drug.