

# Ractopamine Medicated Finishing Cattle Feed Weight Gain, Feed Efficiency and Carcass Leanness

## Type C Medicated Feed

For increased rate of weight gain, improved feed efficiency and increased carcass leanness in cattle fed in confinement for slaughter during the last 28 to 42 days on feed.

### ACTIVE DRUG INGREDIENT

Ractopamine hydrochloride..... 9.8 to 24.6 g/ton\*

### GUARANTEED ANALYSIS

Crude Protein, not less than.....	_____ %
Non-Protein Nitrogen (NPN) <sup>1</sup> , not more than.....	_____ %
Crude Fat, not less than.....	_____ %
Crude Fiber, not more than.....	_____ %
Calcium, not less than.....	_____ %
Calcium, not more than.....	_____ %
Phosphorus, not less than.....	_____ %
Salt <sup>2</sup> , not less than.....	_____ %
Salt <sup>2</sup> , not more than.....	_____ %
Sodium <sup>3</sup> , not less than.....	_____ %
Sodium <sup>3</sup> , not more than.....	_____ %
Potassium, not less than.....	_____ %
Vitamin A <sup>2, 4</sup> , not less than.....	_____ I.U./lb

<sup>1</sup>When added.

<sup>2</sup>If added.

<sup>3</sup>Shall be guaranteed only when total sodium exceeds that furnished by the maximum salt guarantee.

<sup>4</sup>Other than precursors of Vitamin A.

### INGREDIENTS

Each ingredient must be named in accordance with the names and definitions adopted by the Association of American Feed Control Officials.

### FEEDING DIRECTIONS

Feed continuously to cattle fed in confinement for slaughter as the sole ration for last 28 to 42 days on feed.

## **WARNING**

The active ingredient, ractopamine hydrochloride, is a beta-adrenergic agonist. Individuals with cardiovascular disease should exercise special caution to avoid exposure. Not for use in humans. Keep out of the reach of children. The ractopamine hydrochloride formulation (Type A Medicated Article) poses a low dust potential under usual conditions of handling and mixing. When mixing and handling ractopamine hydrochloride, use protective clothing, impervious gloves, protective eye wear, and a NIOSH-approved dust mask. Operators should wash thoroughly with soap and water after handling. If accidental eye contact occurs, immediately rinse eyes thoroughly with water. If irritation persists, seek medical attention. The material safety data sheet contains more detailed occupational safety information. To report adverse effects, access medical information, or obtain additional product information, call 1-888-963-8471.

## **CAUTION**

Not for animals intended for breeding.

**MANUFACTURED BY**  
**BLUE BIRD FEED MILL**  
Any Town, USA 12345

**Net weight lb (kg) on bag or bulk**

\* Final printed label on formulated Type C medicated feed must bear a single drug concentration.