

**BLUE BIRD**  
**Nicarbazin + Bacitracin methylene disalicylate**  
**Type C Broiler Chicken Feed – Cocci/NE**  
**Medicated**

As an aid in preventing outbreaks of cecal (*Eimeria tenella*) and intestinal (*E. acervulina*, *E. maxima*, *E. necatrix* and *E. brunetti*) coccidiosis and as an aid in the prevention of necrotic enteritis caused or complicated by *Clostridium* spp. or other organisms susceptible to bacitracin in broiler chickens.

**Active Drug Ingredients**

Nicarbazin.....90.8 – 181.6 g/ton (0.01 – 0.02%)\*  
 Bacitracin methylene disalicylate..... 50 g/ton

**Guaranteed Analysis**

Crude Protein (Min) .....	_____	%
Lysine (Min).....	_____	%
Methionine (Min) .....	_____	%
Crude Fat (Min) .....	_____	%
Crude Fiber (Max) .....	_____	%
Calcium (Min) .....	_____	%
Calcium (Max) .....	_____	%
Phosphorus (Min).....	_____	%
Salt <sup>a</sup> (Min) .....	_____	%
Salt <sup>a</sup> (Max).....	_____	%
Sodium <sup>b</sup> (Min) .....	_____	%
Sodium <sup>b</sup> (Max) .....	_____	%

<sup>a</sup> Guarantee required only when nutrient added except when the feed is intended, represented or serves as a principal source of the nutrient.  
<sup>b</sup> Sodium guarantee required only when total sodium exceeds that furnished by the maximum salt guarantee.

**Ingredients**

Ingredients as defined by AAFCO.

**Feeding Directions**

Feed continuously as the sole ration from the time chicks are placed on litter until past the time when coccidiosis is ordinarily a hazard.

**Caution**

Do not use as a treatment for coccidiosis. Do not feed to laying hens. Do not use in flushing mashers. If losses exceed 0.5 percent in a 2-day period, obtain an accurate diagnosis and follow the instructions of your veterinarian or poultry pathologist. Broilers fed feed medicated with nicarbazin may show reduced heat tolerance when exposed to high temperature and high humidity to which they have not been accustomed and under severe conditions, fatalities may result. An ample supply of drinking water and adequate ventilation will improve the birds' tolerance of heat.

**Warning**

Withdraw 4 days before slaughter for use levels at or below 113.5 g nicarbazin/ton ( $\leq 0.0125\%$ ).\*  
 Withdraw 5 days before slaughter for use levels above 113.5 g nicarbazin/ton ( $> 0.0125\%$ ).\*

**Manufactured by**

Blue Bird Feed Mill  
 City, State, Zip

**NET WEIGHT ON BAG OR BULK**

\*The final printed feed label must bear a single concentration of nicarbazin, and the WARNING withdrawal statement must reflect this use level.