

BLUE BIRD
Nicarbazin + Bacitracin methylene disalicylate
Type C Broiler Chicken Feed – Cocci/ROG/FE
Medicated

As an aid in preventing outbreaks of cecal (*Eimeria tenella*) and intestinal (*E. acervulina*, *E. maxima*, *E. necatrix* and *E. brunetti*) coccidiosis and for increased rate of weight gain and improved feed efficiency in broiler chickens.

Active Drug Ingredients

Nicarbazin.....90.8 – 181.6 g/ton (0.01 – 0.02%)*
 Bacitracin methylene disalicylate..... 4 - 50 g/ton*

Guaranteed Analysis

Crude Protein (Min).....	_____	%
Lysine (Min).....	_____	%
Methionine (Min).....	_____	%
Crude Fat (Min).....	_____	%
Crude Fiber (Max).....	_____	%
Calcium (Min).....	_____	%
Calcium (Max).....	_____	%
Phosphorus (Min).....	_____	%
Salt ^a (Min).....	_____	%
Salt ^a (Max).....	_____	%
Sodium ^b (Min).....	_____	%
Sodium ^b (Max).....	_____	%

^a Guarantee required only when nutrient added except when the feed is intended, represented or serves as a principal source of the nutrient.

^b Sodium guarantee required only when total sodium exceeds that furnished by the maximum salt guarantee.

Ingredients

Ingredients as defined by AAFCO.

Feeding Directions

Feed continuously as the sole ration from the time chicks are placed on litter until past the time when coccidiosis is ordinarily a hazard.

Caution

Do not use as a treatment for coccidiosis. Do not feed to laying hens. Do not use in flushing mashers. If losses exceed 0.5 percent in a 2-day period, obtain an accurate diagnosis and follow the instructions of your veterinarian or poultry pathologist. Broilers fed feed medicated with nicarbazin may show reduced heat tolerance when exposed to high temperature and high humidity to which they have not been accustomed and under severe conditions, fatalities may result. An ample supply of drinking water and adequate ventilation will improve the birds' tolerance of heat.

Warning

Withdraw 4 days before slaughter for use levels at or below 113.5 g nicarbazin/ton ($\leq 0.0125\%$).**
 Withdraw 5 days before slaughter for use levels above 113.5 g nicarbazin/ton ($> 0.0125\%$).**

Manufactured by

Blue Bird Feed Mill
 City, State, Zip

NET WEIGHT ON BAG OR BULK

*The final printed feed label must bear a single concentration of each drug

**The WARNING withdrawal statement must reflect use level of nicarbazin in the feed.