

**Ractopamine Finishing Cattle Feed Concentrate - TD
Type B Medicated Feed
– to be used in preparation of cattle top dress Type C medicated feed**

Do Not Feed Undiluted

IMPORTANT: MUST BE THOROUGHLY MIXED INTO FEED BEFORE USE

For increased rate of weight gain and improved feed efficiency in cattle fed in confinement for slaughter during the last 28 to 42 days on feed.

ACTIVE DRUG INGREDIENT

Ractopamine hydrochloride..... not to exceed 4000 g/ton*

GUARANTEED ANALYSIS

Crude Protein, not less than.....	_____	%
Non-Protein Nitrogen (NPN) ¹ , not more than.....	_____	%
Crude Fat, not less than.....	_____	%
Crude Fiber, not more than.....	_____	%
Calcium, not less than.....	_____	%
Calcium, not more than.....	_____	%
Phosphorus, not less than.....	_____	%
Salt ² , not less than.....	_____	%
Salt ² , not more than.....	_____	%
Sodium ³ , not less than.....	_____	%
Sodium ³ , not more than.....	_____	%
Potassium, not less than.....	_____	%
Vitamin A ^{2,4} , not less than.....	_____	I.U./lb

¹When added.

²If added.

³Shall be guaranteed only when total sodium exceeds that furnished by the maximum salt guarantee.

⁴Other than precursors of Vitamin A.

INGREDIENTS

Each ingredient must be named in accordance with the names and definitions adopted by the Association of American Feed Control Officials.

* Final printed label on formulated Type B medicated feed must bear a single concentration of this drug.

TYPE B MEDICATED FEED MIXING DIRECTIONS:

Thoroughly mix Ractopamine Finishing Cattle Feed Concentrate TD - Type B Medicated Feed into one ton of appropriate feed ingredients to obtain the proper concentration in the Ractopamine Finishing Cattle Feed Concentrate TD - Type C Top Dress Medicated Feed (see table below for examples).

Concentration of Ractopamine in Type B Medicated Feed grams/pound	Pounds of Ractopamine Finishing Cattle Feed Concentrate TD - Type B Medicated Feed To Add to Non-Medicated Feed to Make One Ton of Type C Medicated Feed	Resulting Ractopamine Concentration in Type C Medicated Top Dress Feed ^a grams/ton
1000	400	200
	800	400
	1600	800
2000	200	200
	400	400
	800	800
4000	100	200
	200	400
	400	800

^a Based on 90% Dry Matter Basis.

TYPE C MEDICATED FEED - FEEDING DIRECTIONS:

Feed a minimum of 1.0 lb per head per day of Ractopamine Finishing Cattle Feed Concentrate TD - Type C Medicated Top Dress Feed continuously to cattle fed in confinement for slaughter to provide 70 to 400 mg/head/day for the last 28 to 42 days on feed.

WARNING: The active ingredient, ractopamine hydrochloride, is a beta-adrenergic agonist. Individuals with cardiovascular disease should exercise special caution to avoid exposure. Not for use in humans. Keep out of the reach of children. The ractopamine hydrochloride formulation (Type A Medicated Article) poses a low dust potential under usual conditions of handling and mixing. When mixing and handling ractopamine hydrochloride, use protective clothing, impervious gloves, protective eye wear, and a NIOSH-approved dust mask. Operators should wash thoroughly with soap and water after handling. If accidental eye contact occurs, immediately rinse eyes thoroughly with water. If irritation persists, seek medical attention. The material safety data sheet contains more detailed occupational safety information. To report adverse effects, access medical information, or obtain additional product information, call 1-888-963-8471.

CAUTION: Not for animals intended for breeding.

MANUFACTURED BY

BULE BIRD FEED MILL
Any town, USA 12345

Net Weight lb (kg) on bag or bulk