

OXYTETRACYCLINE
 TYPE B
 BLUE BIRD
 CHICKEN FEED A
 MEDICATED

CAUTION: Federal law restricts medicated feed containing this veterinary feed directive (VFD) drug to use by or on the order of a licensed veterinarian.

ACTIVE DRUG INGREDIENT

Oxytetracycline 20 g/lb

INDICATIONS FOR USE	OXYTETRACYCLINE Concentration in Type C Medicated Feed	lb of Type B/ton of Feed
For chickens for control of infectious synovitis caused by <i>Mycoplasma synoviae</i> ; control of fowl cholera caused by <i>Pasteurella multocida</i> susceptible to oxytetracycline.	100 - 200 g/ton	5 - 10
For chickens for control of chronic respiratory disease (CRD) and air sac infection caused by <i>Mycoplasma gallisepticum</i> and <i>Escherichia coli</i> susceptible to oxytetracycline.	400 g/ton	20

GUARANTEED ANALYSIS

Crude Protein (Min) %
 Lysine (Min) %
 Methionine (Min) %
 Crude Fat (Min) %
 Crude Fiber (Max) %
 Calcium (Min) %
 Calcium (Max) %
 Phosphorus (Min) %
 Salt, if added (Min) %
 Salt, if added (Max) %
 Total Sodium (Min)* %
 Total Sodium (Max)* %

* Minimum and Maximum percentage of total sodium shall be guaranteed only when total Sodium exceeds that furnished by the maximum salt guarantee.

INGREDIENTS

Ingredients as defined by AAFCO

WARNING: DO NOT USE IN LOW-CALCIUM FEED CONTAINING LESS THAN 0.55% DIETARY CALCIUM. USE IN SUCH FEEDS MAY RESULT IN VIOLATIVE RESIDUES. DO NOT ADMINISTER TO CHICKENS PRODUCING EGGS FOR HUMAN CONSUMPTION.

___ lbs (___ kg) NET WEIGHT

BLUE BIRD FEED MILL
 Robin, IN 00000