

POLOXALENE  
TYPE C

BLUE BIRD

MEDICATED CATTLE LIQUID FEED #1

For control of legume (alfalfa, clover) and wheat pasture bloat in cattle.

**ACTIVE DRUG INGREDIENT**

Poloxalene ..... 7.5 g/lb

**GUARANTEED ANALYSIS**

Crude Protein (Min) ..... %  
Crude Fat (Min) ..... %  
Crude Fiber (Max) ..... %  
Phosphorus (Min) ..... %  
Potassium (Min) ..... %  
Vitamin A (Min) ..... IU/lb

**INGREDIENTS**

Poloxalene Type A medicated article (99.5%) ..... 1.65 % w/w  
Ammonium Polyphosphate ..... 2.66  
Phosphoric Acid (75%) ..... 3.37  
Sulfuric Acid ..... 1.00  
Water ..... 10.00  
Molasses ..... qs to 100.00

Vitamins A and D and trace minerals may be added.

**Important:** To maintain proper distribution of Bloat Guard in liquid feed supplements, the viscosity of the medicated liquid supplement must be no less than 275 cps. During storage the medicated liquid may separate out of liquid supplements within 5 days if not agitated. It is essential that all of the stored mixture be thoroughly remixed (for example, by using a recirculating pump) before being transferred to feeders with free-turning lick wheels. Movement of the lick wheels is sufficient to prevent separation in the feeders.

**FEEDING and MANAGEMENT DIRECTIONS**

This medicated feed is effective for the control of legume (alfalfa, clover) and wheat pasture bloat in cattle, when consumed as recommended.

1. Use a covered liquid feeder with 1 free-turning lick wheel per 25 head at a height so that all cattle can conveniently lick the wheel.
2. The location of the liquid feeder is extremely important for adequate consumption. Place feeder where cattle congregate (for example, near drinking water), thus providing ready access to the medicated liquid feed. In large pastures, additional feeders should be placed so that cattle are never more than 500 feet from a feeder while grazing.
3. For adequate protection, it is essential that each animal consumes the total recommended amount of medication daily.
4. If grazing legumes or wheat pasture, use a liquid feed containing 7.5 g of Bloat Guard per lb (1.65% w/w). Each animal must consume the medicated liquid feed at a rate of 0.2 lb per 100 lb of body weight per day for adequate protection. Example: 2 lb of the 1.65% medicated feed for a 1000 lb animal daily.
5. Introduce the feeding of the medicated liquid feed at least 2 – 5 days before pasturing on legume or wheat to accustom the cattle to the medicated liquid feed and to lick wheel feeding. If the medicated liquid feed feeding is interrupted, this 2 – 5 day introductory feeding should be repeated.

**Cold Weather Note:** At temperatures below 50°F, this product will thicken and may appear cloudy, but is not damaged.

**MANUFACTURED BY**  
**BLUE BIRD FEED MILL**  
**ANYTOWN, USA 12345**

Net Weight \_\_\_ lb (\_\_\_ kg)