

OXYTETRACYCLINE  
 TYPE C  
 BLUE BIRD  
 SWINE FEED  
 MEDICATED

**CAUTION: Federal law restricts medicated feed containing this veterinary feed directive (VFD) drug to use by or on the order of a licensed veterinarian.**

**ACTIVE DRUG INGREDIENT**

Oxytetracycline .....400 to 600 g/ton

INDICATIONS FOR USE	OXYTETRACYCLINE AMOUNT
Treatment of bacterial enteritis caused by <i>Escherichia coli</i> and <i>Salmonella choleraesuis</i> susceptible to oxytetracycline and treatment of bacterial pneumonia caused by <i>Pasteurella multocida</i> susceptible to oxytetracycline.	10 mg/lb of body weight daily* Feed continuously for 7-14 days
For breeding swine for control and treatment of Leptospirosis (reducing the incidence of abortion and shedding of leptospirae) caused by <i>Leptospira pomona</i> susceptible to oxytetracycline.	10 mg/lb of body weight daily* Feed continuously for not more than 14 days

\*Consumption rates may vary and the oxytetracycline concentration in the feed should be adjusted based on consumption rates. For example, at the consumption rate of 4% of animal's body weight, feed has to contain 500 g/ton oxytetracycline.

**GUARANTEED ANALYSIS**

Crude Protein (Min) ..... \_\_\_\_\_ %  
 Lysine (Min) ..... \_\_\_\_\_ %  
 Crude Fat (Min) ..... \_\_\_\_\_ %  
 Crude Fiber (Max) ..... \_\_\_\_\_ %  
 Calcium (Min) ..... \_\_\_\_\_ %  
 Calcium (Max) ..... \_\_\_\_\_ %  
 Phosphorus (Min) ..... \_\_\_\_\_ %  
 Salt, if added (Min) ..... \_\_\_\_\_ %  
 Salt, if added (Max) ..... \_\_\_\_\_ %  
 Sodium (Min)\* ..... \_\_\_\_\_ %  
 Sodium (Max)\* ..... \_\_\_\_\_ %  
 Selenium (Min) ..... \_\_\_\_\_ ppm  
 Zinc (Min) ..... \_\_\_\_\_ ppm

\* Minimum and maximum percentage of total sodium shall be guaranteed only when total sodium exceeds that furnished by the maximum salt guarantee.

**INGREDIENTS**

Ingredients as defined by AAFCO.

\_\_\_\_\_ lbs (\_\_\_\_\_ kg) NET WEIGHT

BLUE BIRD FEED MILL  
 Robin, IN 00000