

NEOMYCIN - OXYTETRACYCLINE  
TYPE C  
BLUE BIRD  
CATTLE FEED A  
MEDICATED

**CAUTION: Federal law restricts medicated feed containing this veterinary feed directive (VFD) drug to use by or on the order of a licensed veterinarian.**

For cattle for prevention and treatment of the early stages of shipping fever.

**ACTIVE DRUG INGREDIENT<sup>1</sup>**

Oxytetracycline (from oxytetracycline dihydrate) equivalent to oxytetracycline hydrochloride ..... 100 to 2000 g/ton  
Neomycin Sulfate ..... 100 to 2000 g/ton

<sup>1</sup> In a one to one ratio.

**GUARANTEED ANALYSIS**

Crude Protein (Min)..... %  
Equivalent crude protein from Non-Protein Nitrogen (NPN) when added (max)..... %  
Crude Fat (Min)..... %  
Crude Fiber (Max)..... %  
Calcium (Min)..... %  
Calcium (Max)..... %  
Phosphorus (Min)..... %  
Salt (Min)..... %  
Salt (Max)..... %  
Selenium (Min)..... ppm  
Potassium (Min)..... %  
Vitamin A (Min)..... IU

**INGREDIENTS**

Ingredients as defined by AAFCO.

**FEEDING DIRECTIONS**

Feed 0.5-2.0 g oxytetracycline and 0.5-2.0 g neomycin sulfate/head/day 3-5 days before and after arrival in feedlots. For example, include in a supplement based on consumption of 2 lbs of supplement per head per day containing 100 to 2000 g/ton oxytetracycline and 100 to 2000 g/ton neomycin sulfate.

**WARNING: A WITHDRAWAL PERIOD HAS NOT BEEN ESTABLISHED FOR USE IN PRERUMINATING CALVES. DO NOT USE IN CALVES TO BE PROCESSED FOR VEAL. A MILK DISCARD TIME HAS NOT BEEN ESTABLISHED FOR USE IN LACTATING DAIRY CATTLE. DO NOT USE IN FEMALE DAIRY CATTLE 20 MONTHS OF AGE OR OLDER. USE OF MORE THAN ONE PRODUCT CONTAINING NEOMYCIN OR FAILURE TO FOLLOW WITHDRAWAL TIMES MAY RESULT IN ILLEGAL DRUG RESIDUES.**

\_\_\_\_ lbs (\_\_\_\_ kg) NET WEIGHT

BLUE BIRD FEED MILL  
Robin, IN 00000