

NEOMYCIN - OXYTETRACYCLINE
 TYPE C
 BLUE BIRD
 CATTLE MILK REPLACER A
 MEDICATED

CAUTION: Federal law restricts medicated feed containing this veterinary feed directive (VFD) drug to use by or on the order of a licensed veterinarian.

ACTIVE DRUG INGREDIENT¹

Oxytetracycline (from oxytetracycline dihydrate) equivalent to oxytetracycline hydrochloride1500 to 2500 g/ton
 Neomycin Sulfate1500 to 2500 g/ton

¹ In a one to one ratio.

INDICATIONS FOR USE	OXYTETRACYCLINE and NEOMYCIN SULFATE AMOUNT
For calves (up to 250 lb) for treatment of bacterial enteritis caused by <i>Escherichia coli</i> susceptible to oxytetracycline; treatment and control of colibacillosis (bacterial enteritis) caused by <i>E. coli</i> susceptible to neomycin.	10 mg/lb of body weight daily* Feed continuously for 7-14 days
For calves and beef and non-lactating dairy cattle for treatment of bacterial enteritis caused by <i>Escherichia coli</i> and bacterial pneumonia (shipping fever complex) caused by <i>Pasteurella multocida</i> susceptible to oxytetracycline; treatment and control of colibacillosis (bacterial enteritis) caused by <i>E. coli</i> susceptible to neomycin.	10 mg/lb of body weight daily* Feed continuously for 7-14 days

*If a calf weighs 100 lbs, consuming 1 gallon of milk replacer per day; each gallon of liquid replacer should contain 1 lb dry milk replacer (containing 2000 g/ton oxytetracycline and 2000 g/ton neomycin sulfate)

GUARANTEED ANALYSIS

Crude Protein (Min)..... _____ %
 Equivalent crude protein from Non-Protein Nitrogen (NPN) when added (max)..... _____ %
 Crude Fat (Min)..... _____ %
 Crude Fiber (Max)..... _____ %
 Calcium (Min)..... _____ %
 Calcium (Max)..... _____ %
 Phosphorus (Min)..... _____ %
 Salt (Min)..... _____ %
 Salt (Max)..... _____ %
 Selenium (Min)..... _____ ppm
 Potassium (Min)..... _____ %
 Vitamin A (Min)..... _____ IU

INGREDIENTS

Ingredients as defined by AAFCO.

CAUTION: TREATMENT SHOULD CONTINUE 24 TO 48 HOURS BEYOND REMISSION OF DISEASE SYMPTOMS.

WARNING: 5-DAY WITHDRAWAL. A WITHDRAWAL PERIOD HAS NOT BEEN ESTABLISHED FOR USE IN PRERUMINATING CALVES. DO NOT USE IN CALVES TO BE PROCESSED FOR VEAL. A MILK DISCARD TIME HAS NOT BEEN ESTABLISHED FOR USE IN LACTATING DAIRY CATTLE. DO NOT USE IN FEMALE DAIRY CATTLE 20 MONTHS OF AGE OR OLDER. USE OF MORE THAN ONE PRODUCT CONTAINING NEOMYCIN OR FAILURE TO FOLLOW WITHDRAWAL TIMES MAY RESULT IN ILLEGAL DRUG RESIDUES.

_____ lbs (_____ kg) NET WEIGHT

BLUE BIRD FEED MILL
 Robin, IN 00000