

NEOMYCIN - OXYTETRACYCLINE
TYPE C
BLUE BIRD
CHICKEN FEED B
MEDICATED

CAUTION: Federal law restricts medicated feed containing this veterinary feed directive (VFD) drug to use by or on the order of a licensed veterinarian.

For chickens for reduction of mortality due to air sacculitis (air sac infection) caused by *Escherichia coli* susceptible to oxytetracycline.

ACTIVE DRUG INGREDIENT¹

Oxytetracycline (from oxytetracycline dihydrate) equivalent to oxytetracycline hydrochloride..... 500 g/ton
Neomycin Sulfate..... 500 g/ton

¹ In a one to one ratio.

GUARANTEED ANALYSIS

Crude Protein (Min)..... %
Crude Lysine (Min) %
Methionine (Min) %
Crude Fat (Min)..... %
Crude Fiber (Min)..... %
Calcium (Min) %
Calcium (Max)..... %
Phosphorus (Min) %
Salt, if added (Min)..... %
Salt, if added (Max) %
Total Sodium (Min) %
Total Sodium (Max) %

(Minimum and maximum percentage of total sodium shall be guaranteed only when total sodium exceeds that furnished by the maximum salt guarantee)

INGREDIENTS

Ingredients as defined by AAFCO

FEEDING DIRECTIONS

Feed 500 g oxytetracycline and 500 g neomycin sulfate per ton of feed continuously for 5 days.

WARNING: WITHDRAW 24 HOURS BEFORE SLAUGHTER. IN LOW CALCIUM FEEDS WITHDRAW 3 DAYS BEFORE SLAUGHTER. DO NOT FEED TO CHICKENS PRODUCING EGGS FOR HUMAN CONSUMPTION.

_____lbs (____kg) NET WEIGHT

BLUE BIRD FEED MILL
Robin, IN 00000