

NEOMYCIN - OXYTETRACYCLINE
 TYPE C
 BLUE BIRD
 SHEEP FEED A
 MEDICATED

CAUTION: Federal law restricts medicated feed containing this veterinary feed directive (VFD) drug to use by or on the order of a licensed veterinarian.

ACTIVE DRUG INGREDIENT¹

Oxytetracycline (from oxytetracycline dihydrate) equivalent to oxytetracycline hydrochloride500 to 1200 g/ton
 Neomycin Sulfate.....500 to 1200 g/ton

¹ In a one to one ratio.

INDICATIONS FOR USE	OXYTETRACYCLINE and NEOMYCIN SULFATE AMOUNT
Treatment of bacterial enteritis caused by <i>Escherichia coli</i> and bacterial pneumonia caused by <i>Pasteurella multocida</i> susceptible to oxytetracycline; treatment and control of colibacillosis (bacterial enteritis) caused by <i>E. coli</i> susceptible to neomycin	10 mg/lb of body weight daily* Feed continuously for 7-14 days

*For example, if a lamb weighs 60 lb, consuming 1 lb of supplement per head per day containing 1200 g/ton oxytetracycline and 1200 g/ton neomycin sulfate.

GUARANTEED ANALYSIS

Crude Protein (Min)..... _____ %
 Equivalent crude protein from Non-Protein Nitrogen (NPN) when added (max) _____ %
 Crude Fat (Min)..... _____ %
 Crude Fiber (Max)..... _____ %
 Calcium (Min) _____ %
 Calcium (Max)..... _____ %
 Phosphorus (Min) _____ %
 Salt (Min)..... _____ %
 Salt (Max)..... _____ %
 Copper (Min) _____ ppm
 Selenium (Min)..... _____ ppm
 Vitamin A (Min)..... _____ IU

Minimum and maximum percentage of total sodium shall be guaranteed only when total sodium exceeds that furnished by the maximum salt guarantee.

Minimum copper if added, or if total copper exceeds 20 ppm

INGREDIENTS

Ingredients as defined by AAFCO

WARNING: WITHDRAW 5 DAYS BEFORE SLAUGHTER.

_____lbs (____kg) NET WEIGHT

BLUE BIRD FEED MILL
 Robin, IN 00000