

NEOMYCIN - OXYTETRACYCLINE  
 TYPE B  
 BLUE BIRD  
 TURKEY FEED C  
 MEDICATED

**CAUTION: Federal law restricts medicated feed containing this veterinary feed directive (VFD) drug to use by or on the order of a licensed veterinarian.**

**ACTIVE DRUG INGREDIENT<sup>1</sup>**

Oxytetracycline (from oxytetracycline dihydrate) equivalent to oxytetracycline hydrochloride ..... 10 g/lb  
 Neomycin Sulfate..... 10 g/lb

<sup>1</sup> In a one to one ratio.

INDICATIONS FOR USE	Drug Concentration in Type C Medicated Feed, g/ton	lb of Type B/ton of feed
Control of complicating bacterial organisms associated with bluecomb (transmissible enteritis, coronaviral enteritis) susceptible to oxytetracycline	400 - 1000	40 - 100

**GUARANTEED ANALYSIS**

Crude Protein (Min)..... %  
 Crude Lysine (Min) ..... %  
 Methionine (Min) ..... %  
 Crude Fat (Min)..... %  
 Crude Fiber (Min)..... %  
 Calcium (Min) ..... %  
 Calcium (Max)..... %  
 Phosphorus (Min) ..... %  
 Salt, if added (Min)..... %  
 Salt, if added (Max)..... %  
 Total Sodium (Min)..... %  
 Total Sodium (Max)..... %

(Minimum and maximum percentage of total sodium shall be guaranteed only when total sodium exceeds that furnished by the maximum salt guarantee)

**INGREDIENTS**

Ingredients as defined by AAFCO

**WARNING: WITHDRAW 5 DAYS BEFORE SLAUGHTER. DO NOT FEED TO TURKEYS PRODUCING EGGS FOR HUMAN CONSUMPTION.**

\_\_\_\_ lbs (\_\_\_\_ kg) NET WEIGHT

BLUE BIRD FEED MILL  
 Robin, IN 00000