

NEOMYCIN - OXYTETRACYCLINE  
TYPE C  
BLUE BIRD  
TURKEY FEED B  
MEDICATED

**CAUTION: Federal law restricts medicated feed containing this veterinary feed directive (VFD) drug to use by or on the order of a licensed veterinarian.**

For turkeys for control of infectious synovitis caused by *Mycoplasma synoviae* susceptible to oxytetracycline.

**ACTIVE DRUG INGREDIENT<sup>1</sup>**

Oxytetracycline (from oxytetracycline dihydrate) equivalent to oxytetracycline hydrochloride..... 200 g/ton  
Neomycin Sulfate..... 200 g/ton

<sup>1</sup> In a one to one ratio.

**GUARANTEED ANALYSIS**

Crude Protein (Min)..... %  
Crude Lysine (Min)..... %  
Methionine (Min)..... %  
Crude Fat (Min) ..... %  
Crude Fiber (Min)..... %  
Calcium (Min)..... %  
Calcium (Max)..... %  
Phosphorus (Min)..... %  
Salt, if added (Min)..... %  
Salt, if added (Max) ..... %  
Total Sodium (Min) ..... %  
Total Sodium (Max)..... %

(Minimum and maximum percentage of total sodium shall be guaranteed only when total sodium exceeds that furnished by the maximum salt guarantee)

**INGREDIENTS**

Ingredients as defined by AAFCO

**FEEDING DIRECTIONS**

Feed 200 g oxytetracycline and 200 g neomycin sulfate per ton of feed continuously for 7-14 days.

**WARNING: WITHDRAW 5 DAYS BEFORE SLAUGHTER. DO NOT FEED TO TURKEYS PRODUCING EGGS FOR HUMAN CONSUMPTION.**

\_\_\_\_ lbs (\_\_\_\_ kg) NET WEIGHT

BLUE BIRD FEED MILL  
Robin, IN 00000