

OXYTETRACYCLINE
 TYPE C
 BLUE BIRD
 SWINE FEED
 MEDICATED

CAUTION: Federal law restricts medicated feed containing this veterinary feed directive (VFD) drug to use by or on the order of a licensed veterinarian.

ACTIVE DRUG INGREDIENT

Oxytetracycline400 to 600 g/ton

INDICATIONS FOR USE	OXYTETRACYCLINE AMOUNT
Treatment of bacterial enteritis caused by <i>Escherichia coli</i> and <i>Salmonella choleraesuis</i> susceptible to oxytetracycline and treatment of bacterial pneumonia caused by <i>Pasteurella multocida</i> susceptible to oxytetracycline.	10 mg/lb of body weight daily* Feed continuously for 7-14 days
For breeding swine for control and treatment of Leptospirosis (reducing the incidence of abortion and shedding of leptospirae) caused by <i>Leptospira pomona</i> susceptible to oxytetracycline.	10 mg/lb of body weight daily* Feed continuously for not more than 14 days

*Consumption rates may vary and the oxytetracycline concentration in the feed should be adjusted based on consumption rates. For example, at the consumption rate of 4% of animal's body weight, feed has to contain 500 g/ton oxytetracycline.

GUARANTEED ANALYSIS

Crude Protein (Min) %
 Lysine (Min) %
 Crude Fat (Min) %
 Crude Fiber (Max) %
 Calcium (Min) %
 Calcium (Max) %
 Phosphorus (Min) %
 Salt, if added (Min) %
 Salt, if added (Max) %
 Sodium (Min)* %
 Sodium (Max)* %
 Selenium (Min) ppm
 Zinc (Min) ppm

* Minimum and maximum percentage of total sodium shall be guaranteed only when total sodium exceeds that furnished by the maximum salt guarantee.

INGREDIENTS

Ingredients as defined by AAFCO.

____ lbs (____ kg) NET WEIGHT

BLUE BIRD FEED MILL
 Robin, IN 00000