

NEOMYCIN - OXYTETRACYCLINE
TYPE C
BLUE BIRD
TURKEY FEED C
MEDICATED

CAUTION: Federal law restricts medicated feed containing this veterinary feed directive (VFD) drug to use by or on the order of a licensed veterinarian.

For turkeys for control of complicating bacterial organisms associated with bluecomb (transmissible enteritis, coronaviral enteritis) susceptible to oxytetracycline.

ACTIVE DRUG INGREDIENT¹

Oxytetracycline (from oxytetracycline dihydrate) equivalent to oxytetracycline hydrochloride..... 400 to 1000 g/ton
Neomycin Sulfate..... 400 to 1000 g/ton

¹ In a one to one ratio.

GUARANTEED ANALYSIS

Crude Protein (Min).....	_____ %
Crude Lysine (Min).....	_____ %
Methionine (Min).....	_____ %
Crude Fat (Min).....	_____ %
Crude Fiber (Min).....	_____ %
Calcium (Min).....	_____ %
Calcium (Max).....	_____ %
Phosphorus (Min).....	_____ %
Salt, if added (Min).....	_____ %
Salt, if added (Max).....	_____ %
Total Sodium (Min).....	_____ %
Total Sodium (Max).....	_____ %

(Minimum and maximum percentage of total sodium shall be guaranteed only when total sodium exceeds that furnished by the maximum salt guarantee)

INGREDIENTS

Ingredients as defined by AAFCO

FEEDING DIRECTIONS

Feed 25 mg oxytetracycline and 25 mg neomycin sulfate per pound of body weight daily continuously for 7-14 days. Consumption rates may vary and the oxytetracycline and neomycin sulfate concentrations in the feed should be adjusted based on consumption rates. For example, if a bird weighs 15 lb, consuming 0.75 lb of complete feed per day, the feed should contain 1000 g/ton oxytetracycline and 1000 g/ton neomycin sulfate.

WARNING: WITHDRAW 5 DAYS BEFORE SLAUGHTER. DO NOT FEED TO TURKEYS PRODUCING EGGS FOR HUMAN CONSUMPTION.

_____ lbs (_____ kg) NET WEIGHT

BLUE BIRD FEED MILL
Robin, IN 00000