

NR 44 v 12010 5-0137

PYRANTEL TARTRATE
TYPE B MEDICATED SWINE FEED

BLUE BIRD

Do not feed undiluted.

For the removal and control of large roundworm (*Ascaris suum*) and nodular worm (*Oesophagostomum*) infections in swine. To aid in the prevention of migration and establishment of large roundworm (*Ascaris suum*) infections; aid in the prevention of establishment of nodular worm (*Oesophagostomum*) infections in swine. For the removal and control of large roundworm (*Ascaris suum*) infections in swine.

ACTIVE DRUG INGREDIENT

Pyrantel tartrate4.8 g/lb

GUARANTEED ANALYSIS

Crude Protein (Min)	_____	%
Lysine (Min).....	_____	%
Crude Fat (Min).....	_____	%
Crude Fiber (Max).....	_____	%
Calcium (Min).....	_____	%
Calcium (Max)	_____	%
Phosphorus (Min).....	_____	%
Salt (Min)	_____	%
Salt (Max).....	_____	%
Selenium.....	_____	%
Zinc (Min)	_____	%

(Minimum and maximum percentage of total Sodium shall be guaranteed only when total Sodium exceeds that furnished by the maximum salt guarantee)

INGREDIENTS

Ingredients as defined by AAFCO

MIXING DIRECTIONS

INDICATION FOR USE	PYRANTEL TARTRATE CONCENTRATION	lb of Type B/ton of feed
For the removal and control of large roundworm (<i>Ascaris suum</i>) and nodular worm (<i>Oesophagostomum</i>) infections in swine	800 g/ton	166.7 lb
Aid in the prevention of migration and establishment of large roundworm (<i>Ascaris suum</i>) infections; aid in the prevention of establishment of nodular worm (<i>Oesophagostomum</i>) infections in swine.	96 g/ton	20 lb
For the removal and control of large roundworm (<i>Ascaris suum</i>) infections in swine.	96 g/ton	20 lb

CAUTION: Consult your veterinarian before using in severely debilitated animals. Do not mix in feeds containing bentonite.

WARNING: Withdraw 24 hours prior to slaughter.

BLUE BIRD FEED MILL
ROBIN, IN 00000

NET WEIGHT ON BAG OR BULK