

**PYRANTEL TARTRATE
TYPE C MEDICATED SWINE FEED
BLUE BIRD**

For the removal and control of large roundworm (*Ascaris suum*) and nodular worm (*Oesophagostomum*) infections in swine. To aid in the prevention of migration and establishment of large roundworm (*Ascaris suum*) infections; aid in the prevention of establishment of nodular worm (*Oesophagostomum*) infections in swine. For the removal and control of large roundworm (*Ascaris suum*) infections in swine.

ACTIVE DRUG INGREDIENT

Pyrantel tartrate 96 or 800 g/ton*

GUARANTEED ANALYSIS

Crude Protein (Min)	_____ %
Lysine (Min).....	_____ %
Crude Fat (Min).....	_____ %
Crude Fiber (Max).....	_____ %
Calcium (Min).....	_____ %
Calcium (Max).....	_____ %
Phosphorus (Min).....	_____ %
Salt (Min).....	_____ %
Salt (Max).....	_____ %
Selenium.....	_____ %
Zinc (Min).....	_____ %

(Minimum and maximum percentage of total Sodium shall be guaranteed only when total Sodium exceeds that furnished by the maximum salt guarantee)

INGREDIENTS

Ingredients as defined by AAFCO

FEEDING DIRECTIONS

INDICATIONS FOR USE		PYRANTEL TARTRATE CONCENTRATION			DIRECTIONS FOR USE		
For the removal and control of large roundworm (<i>A. suum</i>) and nodular worm (<i>Oesophagostomum</i>) infections in swine		800 g/ton			As a single therapeutic treatment in complete feed; feed at the rate of feed per 40 lb of body weight for animals up to 200 lb. For animals 200 lb or over, feed 5 lb feed per head. Quantity of medicated feed required to treat pigs of various body weight is show below:		
Pig Weight (lb)	20	40	60	80	100	200 or over	
Quantity of Feed per Pig	½ lb	1 lb	1 ½ lb	2 lb	2 ½ lb	5 lb	
Fast pigs overnight for optimum results. Pigs will generally consume this quantity of medicated feed within a few hours time after which regular feeding should be resumed. Better worming is achieved when pigs are separated by size into different lots or pens for treatment. Water should be made available to animals during fasting and treatment periods							
Aid in the prevention of migration and establishment of large roundworm (<i>A. suum</i>) infections; aid in the prevention of establishment of nodular worm (<i>Oesophagostomum</i>) infections in swine.		96 g/ton			Feed continuously as the sole ration.		
For the removal and control of large roundworm (<i>Ascaris suum</i>) infections in swine.		96 g/ton			Feed for 3 days as the sole ration.		

CAUTION: Consult your veterinarian before using in severely debilitated animals. Do not mix in feeds containing bentonite.

WARNING: Withdraw 24 hours prior to slaughter.

BLUE BIRD FEED MILL
ROBIN, IN 00000

NET WEIGHT ON BAG OR BULK

* One of these concentrations should be specified.