

**THIS BLUEBIRD LABEL IS INTENDED FOR USE ONLY WITH
Zilpaterol Finishing Cattle Feed
+ Monensin and Tylosin Type C Medicated Component Feed**

**HEIFER TYPE C MEDICATED FEED – BAG or BULK
For Beef and Dairy Heifers**

Caution: Federal law restricts medicated feed containing this veterinary feed directive (VFD) drug to use by or on the order of a licensed veterinarian.

Heifers Fed in Confinement for Slaughter: For increased rate of weight gain, improved feed efficiency, and suppression of estrus (heat).

ACTIVE DRUG INGREDIENT

Melengestrol acetate..... (0.0000276-0.00022%)*
Each pound contains (0.125-1.0) mg melengestrol acetate (select one)*

GUARANTEED ANALYSIS

Crude Protein (Min)₁..... %
NPN (Max)..... %
Crude Fat (Min)..... %
Crude Fiber (Max)..... %
Calcium (Min)..... %
Calcium (Max)..... %
Phosphorus (Min)..... %
Salt₂ (Min)..... %
Salt₂ (Max)..... %
Sodium₃ (Min)..... %
Sodium₃ (Max)..... %
Potassium (Min)..... %
Vitamin A_{2,4} (Min)..... I.U./lb

¹ When added.

² If added.

³ Shall be guaranteed only when total sodium exceeds that furnished by the maximum salt guarantee.

⁴ Other than precursors of Vitamin A.

INGREDIENTS

Each ingredient must be named in accordance with the names and definitions adopted by the Association of American Feed Control Officials.

*Final printed label of Type C Medicated Feed must bear a single drug concentration of this drug.

FEEDING DIRECTIONS

Must be top dressed or mixed with a Type C feed containing 6.8 to 24 g/ton zilpaterol (to provide 60 mg/head/day via component feed), 10 to 40 g/ton monensin, and 8-10 g/ton tylosin.

Feed at the rate of 0.5 to 2.0 lb/head/day (specify one level) to provide 0.25 to 0.5 mg melengestrol acetate/head/day (specify one level). Feed melengestrol acetate in this combination for the final 20 to 40 days on feed.

CAUTION: Melengestrol acetate is not effective in steers and spayed heifers. Withdrawal periods of three to five days or more should be avoided to prevent the possibility that the heifers may come into estrus (heat) at loading time.

Do not use in any finished feed (supplement, concentrate or complete feed) containing in excess of 2% bentonite.

MANUFACTURED BY
BLUE BIRD FEED MILL
Any town, USA 12345

Net Weight lb (kg) on bag or bulk

Batch (Lot or Control) No.:

NADA 141-280 Apr 2016