

CHLORTETRACYCLINE
TYPE C
BLUE BIRD
CATTLE MILK REPLACER
MEDICATED

CAUTION: Federal law restricts medicated feed containing this veterinary feed directive (VFD) drug to use by or on the order of a licensed veterinarian.

For calves (up to 250 lb) for the treatment of bacterial enteritis caused by *E coli* susceptible to chlortetracycline.

ACTIVE DRUG INGREDIENT

Chlortetracycline 2000 g/ton

GUARANTEED ANALYSIS

Crude Protein (Min).....	_____%
Non-Protein Nitrogen (NPN, max) ^a	_____%
Crude Fat (min).....	_____%
Crude Fiber (max).....	_____%
Calcium (min) ^a	_____%
Calcium (max) ^a	_____%
Phosphorus (min) ^a	_____%
Salt (min) ^a	_____%
Salt (max) ^a	_____%
Sodium (min) ^b	_____%
Sodium (max) ^b	_____%
Selenium (min).....	_____ppm
Potassium (min) ^a	_____%
Vitamin A (min) ^a	_____IU

^a Required only when nutrient added.

^b Required only when total sodium exceeds that furnished by the maximum salt guarantee.

INGREDIENTS

Ingredients as defined by AAFCO

FEEDING DIRECTIONS

Feed 10 mg chlortetracycline/lb of body weight daily. For example, for a calf weighing 100 lb, consuming 1 lb of dry milk replacer per day mixed with 1 gallon of water, the milk replacer should contain 2000 g chlortetracycline/ton. Limit use to 5 days.

WARNING: AT 10 MG/LB USE LEVEL, WITHDRAW 10 DAYS BEFORE SLAUGHTER. A WITHDRAWAL PERIOD HAS NOT BEEN ESTABLISHED IN PRERUMINATING CALVES. DO NOT USE IN CALVES TO BE PROCESSED FOR VEAL.

BLUE BIRD FEED MILL
Robin, IN 00000

NET WEIGHT on Bag or Bulk