

NEOMYCIN - OXYTETRACYCLINE  
 TYPE B  
 BLUE BIRD  
 CATTLE FEED B  
 MEDICATED

**CAUTION: Federal law restricts medicated feed containing this veterinary feed directive (VFD) drug to use by or on the order of a licensed veterinarian.**

**ACTIVE DRUG INGREDIENT<sup>1</sup>**

Oxytetracycline (from oxytetracycline dihydrate) equivalent to oxytetracycline hydrochloride ..... 10 g/lb  
 Neomycin Sulfate..... 10 g/lb

<sup>1</sup> In a one to one ratio.

| INDICATIONS FOR USE   | Drug Concentration in Type C medicated feed, g/ton | lb of Type B/ton of Type C Feed |
|---|--|---------------------------------|
| For calves and beef and non-lactating dairy cattle for treatment of bacterial enteritis caused by <i>Escherichia coli</i> and bacterial pneumonia (shipping fever complex) caused by <i>Pasteurella multocida</i> susceptible to oxytetracycline; treatment and control of colibacillosis (bacterial enteritis) caused by <i>E. coli</i> susceptible to neomycin. | 1000 - 5000  | 100 - 500                       |

**GUARANTEED ANALYSIS**

Crude Protein (Min)..... %  
 Equivalent crude protein from Non-Protein Nitrogen (NPN) when added (max)..... %  
 Crude Fat (Min) ..... %  
 Crude Fiber (Max) ..... %  
 Calcium (Min) ..... %  
 Calcium (Max)..... %  
 Phosphorus (Min) ..... %  
 Salt (Min)..... %  
 Salt (Max)..... %  
 Selenium (Min)..... %  
 Potassium (Min)..... %  
 Vitamin A (Min)..... %

**INGREDIENTS**

Ingredients as defined by AAFCO.

**WARNING: A WITHDRAWAL PERIOD HAS NOT BEEN ESTABLISHED FOR USE IN PRERUMINATING CALVES. DO NOT USE IN CALVES TO BE PROCESSED FOR VEAL. A MILK DISCARD TIME HAS NOT BEEN ESTABLISHED FOR USE IN LACTATING DAIRY CATTLE. DO NOT USE IN FEMALE DAIRY CATTLE 20 MONTHS OF AGE OR OLDER. WITHDRAW 5 DAYS BEFORE SLAUGHTER. USE OF MORE THAN ONE PRODUCT CONTAINING NEOMYCIN OR FAILURE TO FOLLOW WITHDRAWAL TIMES MAY RESULT IN ILLEGAL DRUG RESIDUES.**

\_\_\_\_lbs (\_\_\_\_kg) NET WEIGHT

BLUE BIRD FEED MILL  
 Robin, IN 00000