

NEOMYCIN - OXYTETRACYCLINE
TYPE C

BLUE BIRD
CATTLE FEED C
MEDICATED

CAUTION: Federal law restricts medicated feed containing this veterinary feed directive (VFD) drug to use by or on the order of a licensed veterinarian.

ACTIVE DRUG INGREDIENT¹

Oxytetracycline (from oxytetracycline dihydrate) equivalent to oxytetracycline hydrochloride 5 to 75 g/ton
Neomycin Sulfate 5 to 75 g/ton

¹ In a one to one ratio.

INDICATIONS FOR USE	OXYTETRACYCLINE and NEOMYCIN SULFATE AMOUNT	DRUG CONCENTRATIONS IN TYPE C MEDICATED FEEDS, g/ton
For growing cattle (over 400 lb) for reduction of liver condemnation due to liver abscesses	75 mg/head/day ¹ Feed continuously	15 - 75

¹If a calf weighs 500 lb, consuming 2 lb of supplement per head per day containing 75 g/ton oxytetracycline and 75 g/ton neomycin sulfate

GUARANTEED ANALYSIS

Crude Protein (Min)..... _____ %
 Equivalent crude protein from Non-Protein Nitrogen (NPN) when added (max)..... _____ %
 Crude Fat (Min) _____ %
 Crude Fiber (Max) _____ %
 Calcium (Min) _____ %
 Calcium (Max)..... _____ %
 Phosphorus (Min) _____ %
 Salt (Min)..... _____ %
 Salt (Max)..... _____ %
 Selenium (Min)..... _____ ppm
 Potassium (Min)..... _____ %
 Vitamin A (Min) _____ IU

INGREDIENTS

Ingredients as defined by AAFCO.

WARNING: A WITHDRAWAL PERIOD HAS NOT BEEN ESTABLISHED IN PRERUMINATING CALVES. DO NOT USE IN CALVES TO BE PROCESSED FOR VEAL. USE OF MORE THAN ONE PRODUCT CONTAINING NEOMYCIN OR FAILURE TO FOLLOW WITHDRAWAL TIMES MAY RESULT IN ILLEGAL DRUG RESIDUES.

_____ lbs (_____ kg) NET WEIGHT

BLUE BIRD FEED MILL
Robin, IN 00000