

**THIS BLUEBIRD LABEL IS INTENDED FOR USE ONLY WITH  
Zilpaterol, Monensin, and Tylosin Finishing Cattle Feed  
Type C Medicated Complete Feed**

**HEIFER TYPE C MEDICATED LIQUID FEED – BAG or BULK  
For Beef and Dairy Heifers**

**Caution:** Federal law restricts medicated feed containing this veterinary feed directive (VFD) drug to use by or on the order of a licensed veterinarian.

Heifers Fed in Confinement for Slaughter: For increased rate of weight gain, improved feed efficiency, and suppression of estrus (heat).

**ACTIVE DRUG INGREDIENT**

Melengestrol acetate..... (0.0000276-0.00022%)\*  
Each pound contains (0.125-1.0) mg melengestrol acetate ..... (select one)\*

**GUARANTEED ANALYSIS**

Crude Protein (Min)<sub>1</sub>..... %  
NPN (Max)..... %  
Crude Fat (Min)..... %  
Crude Fiber (Max)..... %  
Calcium (Min)..... %  
Calcium (Max)..... %  
Phosphorus (Min)..... %  
Salt<sub>2</sub> (Min)..... %  
Salt<sub>2</sub> (Max)..... %  
Sodium<sub>3</sub> (Min)..... %  
Sodium<sub>3</sub> (Max)..... %  
Potassium (Min)..... %  
Vitamin A<sub>2,4</sub> (Min)..... I.U./lb

<sup>1</sup> When added.

<sup>2</sup> If added.

<sup>3</sup> Shall be guaranteed only when total sodium exceeds that furnished by the maximum salt guarantee.

<sup>4</sup> Other than precursors of Vitamin A.

**INGREDIENTS**

Each ingredient must be named in accordance with the names and definitions adopted by the Association of American Feed Control Officials.

---

\*Final printed label of Type C Medicated Feed must bear a single drug concentration of this drug.

## **FEEDING DIRECTIONS**

Must be top dressed or mixed with a Type C feed containing 6.8 g/ton zilpaterol, 10 to 40 g/ton monensin, and 8-10 g/ton tylosin.

Feed at the rate of 0.5 to 2.0 lb/head/day (specify one level) to provide 0.25 to 0.5 mg melengestrol acetate/head/day (specify one level). Feed melengestrol acetate in this combination for the final 20 to 40 days on feed.

## **MIXING DIRECTIONS**

Mixing directions for liquid Type C feeds stored in recirculation tank systems are: "Recirculate immediately prior to use for no less than 10 minutes, moving not less than 1 percent of the tank contents from the bottom of the tank to the top. Recirculate daily, as directed in this paragraph even when the Type C feed is not used." Mixing directions for liquid Type C feeds stored in mechanical, air or other agitation-type tank systems are: "Agitate immediately prior to use for not less than 10 minutes, creating a turbulence at the bottom of the tank that is visible at the top. Agitate daily, as directed in this paragraph, even when the Type C feed is not used."

**CAUTION:** Melengestrol acetate is not effective in steers and spayed heifers. Withdrawal periods of three to five days or more should be avoided to prevent the possibility that the heifers may come into estrus (heat) at loading time.

Do not use in any finished feed (supplement, concentrate or complete feed) containing in excess of 2% bentonite.

**MANUFACTURED BY**  
BLUE BIRD FEED MILL  
Any town, USA 12345

**Net Weight lb (kg) on bag or bulk**

**Batch (Lot or Control) No.:**

(NADA 141-280) Apr 2016