

OXYTETRACYCLINE – SALINOMYCIN SODIUM
 TYPE C
 BLUE BIRD
 CHICKEN FEED
 MEDICATED

CAUTION: Federal law restricts medicated feed containing this veterinary feed directive (VFD) drug to use by or on the order of a licensed veterinarian.

For chickens as an aid in the reduction of mortality due to air-sacculitis (air sac infection) caused by *Esherichia coli* sensitive to oxytetracycline and for the prevention of coccidiosis caused by *Eimeria necatrix*, *E. tenella*, *E. acervulina*, *E. brunetti*, *E. mivati* and *E. maxima*.

ACTIVE DRUG INGREDIENT

Oxytetracycline 500 g/ton
 Salinomycin Sodium 40 - 60 g/ton*

GUARANTEED ANALYSIS

Crude Protein (Min)..... %
 Crude Lysine (Min) %
 Methionine (Min) %
 Crude Fat (Min)..... %
 Crude Fiber (Min)..... %
 Calcium (Min) %
 Calcium (Max)..... %
 Phosphorus (Min) %
 Salt, if added (Min)..... %
 Salt, if added (Max)..... %
 Total Sodium (Min) %
 Total Sodium (Max) %

(Minimum and maximum percentage of total sodium shall be guaranteed only when total sodium exceeds that furnished by the maximum salt guarantee)

INGREDIENTS

Ingredients as defined by AAFCO

FEEDING DIRECTIONS

Feed for 5 days as sole ration. Treat at first clinical signs of disease.

WARNING: DO NOT FEED TO LAYING CHICKENS. DO NOT USE IN LOW-CALCIUM FEED CONTAINING LESS THAN 0.55% DIETARY CALCIUM. USE IN SUCH FEEDS MAY RESULT IN VIOLATIVE RESIDUES. WITHDRAW 24 HOURS BEFORE SLAUGHTER.

CAUTION: Do not use in feeds containing pellet binders. May be fatal if accidentally fed to adult turkeys or to horses.

____ lbs (____ kg) NET WEIGHT

BLUE BIRD FEED MILL
 Robin, IN 00000

* Final printed label on formulated Type C medicated feed must bear a single concentration of drug.