

Amprolium and Bambermycins Broiler Chickens

Type C Medicated Feed

Prevention of coccidiosis where immunity to coccidiosis is not desired and for increased rate of weight gain and improved feed efficiency in broiler chickens.

ACTIVE DRUG INGREDIENT

Amprolium 113.5 to 227 g/ton (0.0125 to 0.025%)*
 Bambermycins..... 1 to 2 g/ton*

GUARANTEED ANALYSIS

Crude Protein, not less than.....	_____	%
Lysine, not less than.....	_____	%
Methionine, not less than.....	_____	%
Crude Fat, not less than.....	_____	%
Crude Fiber, not more than.....	_____	%
Calcium, not less than.....	_____	%
Calcium, not more than.....	_____	%
Phosphorus, not less than.....	_____	%
Salt ¹ , not less than.....	_____	%
Salt ¹ , not more than.....	_____	%
Sodium ² , not less than.....	_____	%
Sodium ² , not more than.....	_____	%

¹ If added.

² Shall be guaranteed only when total sodium exceeds that furnished by the maximum salt guarantee.

INGREDIENTS

Each ingredient must be named in accordance with the names and definitions adopted by the Association of American Feed Control Officials.

DIRECTIONS FOR USE

Feed continuously as the sole ration. Increasing levels of amprolium may not prevent coccidiosis caused by strains of *Eimeria* commonly found to be resistant to amprolium. Use 0.0125% amprolium for most field conditions as they exist under modern management practices. Where severe coccidiosis conditions exist and where immunity is not required, use 0.025% amprolium.

CAUTION: Use as sole source of amprolium. Do not use in feeds containing bentonite. Do not change the litter while giving this feed unless absolutely necessary. If losses exceed 0.5% in a 2-day period, obtain an accurate diagnosis and follow the instructions of your veterinarian or poultry pathologist. Losses may result from intercurrent disease or other conditions affecting drug intake, which can contribute to the virulence of coccidiosis under field conditions.

MANUFACTURED BY

BLUE BIRD FEED MILL
 Anytown, USA 12345

Net weight lb (kg) on bag or bulk

*Final printed label on formulated Type C medicated feed must bear a single concentration of each drug.