

## Clopidol and Bambermycins WD5 Broiler Chickens

### Type C Medicated Feed

As an aid in the prevention of coccidiosis caused by *Eimeria tenella*, *E. necatrix*, *E. acervulina*, *E. maxima*, *E. brunetti*, and *E. mivati* and for increased rate of weight gain and improved feed efficiency in broiler chickens.

#### ACTIVE DRUG INGREDIENT

Clopidol.....	227 g/ton
Bambermycins .....	1 to 2 g/ton*

#### GUARANTEED ANALYSIS

Crude Protein, not less than .....	_____ %
Lysine, not less than.....	_____ %
Methionine, not less than.....	_____ %
Crude Fat, not less than .....	_____ %
Crude Fiber, not more than.....	_____ %
Calcium, not less than .....	_____ %
Calcium, not more than .....	_____ %
Phosphorus, not less than .....	_____ %
Salt <sup>1</sup> , not less than .....	_____ %
Salt <sup>1</sup> , not more than.....	_____ %
Sodium <sup>2</sup> , not less than.....	_____ %
Sodium <sup>2</sup> , not more than.....	_____ %

<sup>1</sup> If added.

<sup>2</sup> Shall be guaranteed only when total sodium exceeds that furnished by the maximum salt guarantee.

#### INGREDIENTS

Each ingredient must be named in accordance with the names and definitions adopted by the Association of American Feed Control Officials.

#### DIRECTIONS FOR USE

Feed continuously as the sole ration until 5 days before slaughter.

**WARNING:** Withdraw 5 days before slaughter or feed 113.5 g/ton clopidol and 1-2 g/ton bambermycins during those 5 days before slaughter. Do not feed to chickens over 16 weeks of age.

**CAUTION:** Consult a veterinarian or poultry pathologist if losses exceed 0.5% in a 2-day period.

#### MANUFACTURED BY

**BLUE BIRD FEED MILL**  
Anytown, USA 12345

Net weight lb (kg) on bag or bulk

\*Final printed label on formulated Type C medicated feed must bear a single concentration of the drug.