

**BLUE BIRD  
BMD  
Type B Chicken/Turkey/Pheasant Feed – ROG/FE  
Medicated**

For increased rate of weight gain and improved feed efficiency in broiler and replacement chickens,  
growing turkeys, and pheasants.

**Active Drug Ingredient**

Bacitracin methylene disalicylate.....51 – 50,000 g/ton\*  
(0.0255 – 25 g/lb)\*

**Guaranteed Analysis**

Crude Protein (Min).....	_____	%
Lysine (Min).....	_____	%
Methionine (Min).....	_____	%
Crude Fat (Min).....	_____	%
Crude Fiber (Max).....	_____	%
Calcium (Min).....	_____	%
Calcium (Max).....	_____	%
Phosphorus (Min).....	_____	%
Salt <sup>a</sup> (Min).....	_____	%
Salt <sup>a</sup> (Max).....	_____	%
Sodium <sup>b</sup> (Min).....	_____	%
Sodium <sup>b</sup> (Max).....	_____	%

<sup>a</sup> Guarantee required only when nutrient added except when the feed is intended, represented or serves as a principal source of the nutrient.

<sup>b</sup> Sodium guarantee required only when total sodium exceeds that furnished by the maximum salt guarantee.

**Ingredients**

Ingredients as defined by AAFCO.

**Mixing Directions**

Thoroughly mix this Type B medicated feed with non-medicated feed ingredients to manufacture one ton of Type C medicated feed for [chickens, turkeys, pheasants] containing 4 – 50 g bacitracin methylene disalicylate per ton.

*The following table provides examples of mixing rates:*

Type B BMD concentration (g/ton)	Type B per ton of Type C (lb)	Non-medicated feed per ton of Type C (lb)	Type C BMD concentration (g/ton)
10,000 (5 g/lb)	0.8	1999.2	4
10,000 (5 g/lb)	5	1995	25
10,000 (5 g/lb)	10	1990	50
50,000 (25 g/lb)	0.2	1999.8	5
50,000 (25 g/lb)	2	1998	50

The resulting Type C medicated feed should be fed continuously as the sole ration.

**Manufactured by**  
Blue Bird Feed Mill  
City, State, Zip

**NET WEIGHT ON BAG OR BULK**

\*The final printed feed label should list only a single drug concentration.