

**BLUE BIRD  
BMD  
Type B Turkey Feed – TE  
Medicated**

As an aid in control of transmissible enteritis in growing turkeys complicated by organisms susceptible to bacitracin methylene disalicylate.

**Active Drug Ingredient**

Bacitracin methylene disalicylate.....201 – 50,000 g/ton\*  
(0.10 – 25 g/lb)\*

**Guaranteed Analysis**

Crude Protein (Min).....	_____	%
Lysine (Min) .....	_____	%
Methionine (Min) .....	_____	%
Crude Fat (Min).....	_____	%
Crude Fiber (Max).....	_____	%
Calcium (Min).....	_____	%
Calcium (Max).....	_____	%
Phosphorus (Min) .....	_____	%
Salt <sup>a</sup> (Min) .....	_____	%
Salt <sup>a</sup> (Max) .....	_____	%
Sodium <sup>b</sup> (Min).....	_____	%
Sodium <sup>b</sup> (Max).....	_____	%

<sup>a</sup> Guarantee required only when nutrient added except when the feed is intended, represented or serves as a principal source of the nutrient.

<sup>b</sup> Sodium guarantee required only when total sodium exceeds that furnished by the maximum salt guarantee.

**Ingredients**

Ingredients as defined by AAFCO.

Thoroughly mix this Type B medicated feed with non-medicated feed ingredients to manufacture one ton of Type C medicated feed for turkeys containing 200 g bacitracin methylene disalicylate per ton.

The following table provides examples of mixing rates:

Type B BMD concentration (g/ton)	Type B per ton of Type C (lb)	Non-medicated feed per ton of Type C (lb)	Type C BMD concentration (g/ton)
10,000 (5 g/lb)	40	1960	200
20,000 (10 g/lb)	20	1980	200
50,000 (25 g/lb)	8	1992	200

The resulting Type C medicated feed should be fed continuously as the sole ration.

**Manufactured by**  
Blue Bird Feed Mill  
City, State, Zip

**NET WEIGHT ON BAG OR BULK**

\*The final printed feed label should list only a single drug concentration.