

**BLUE BIRD  
BMD  
Type C Pregnant Sow Feed  
Medicated**

For control of clostridial enteritis caused by *Clostridium perfringens* in suckling piglets when fed to sows from 14 days before through 21 days after farrowing on premises with a history of clostridial scours.

**Active Drug Ingredient**

Bacitracin methylene disalicylate.....250 g/ton

**Guaranteed Analysis**

Crude Protein (Min) .....	_____	%
Lysine (Min) .....	_____	%
Crude Fat (Min) .....	_____	%
Crude Fiber (Max) .....	_____	%
Calcium (Min) .....	_____	%
Calcium (Max) .....	_____	%
Phosphorus (Min).....	_____	%
Salt <sup>a</sup> (Min).....	_____	%
Salt <sup>a</sup> (Max).....	_____	%
Sodium <sup>b</sup> (Min).....	_____	%
Sodium <sup>b</sup> (Max).....	_____	%
Selenium <sup>c</sup> (Min) .....	_____	ppm or %
Zinc (Min) <sup>c</sup> .....	_____	ppm or %

<sup>a</sup> Guarantee required only when nutrient added except when the feed is intended, represented or serves as a principal source of the nutrient.

<sup>b</sup> Sodium guarantee required only when total sodium exceeds that furnished by the maximum salt guarantee.

<sup>c</sup> Guarantee shall be stated in ppm when less than 10,000 ppm and in percentage when concentration is 10,000 ppm (1%) or greater.

**Ingredients**

Ingredients as defined by AAFCO.

**Feeding Directions**

Feed as the sole ration to pregnant sows from 14 days before through 21 days after farrowing.

**Caution**

Diagnosis should be confirmed by a veterinarian when results are not satisfactory.

**Manufactured by**  
Blue Bird Feed Mill  
City, State, Zip

**NET WEIGHT ON BAG OR BULK**