

**THIS BLUEBIRD LABEL IS INTENDED FOR USE ONLY WITH
Zilpaterol Finishing Cattle Feed
Type C Medicated Component Feed**

**HEIFER TYPE C MEDICATED LIQUID FEED – BAG or BULK
For Beef and Dairy Heifers**

Heifers Fed in Confinement for Slaughter: For increased rate of weight gain, improved feed efficiency, and suppression of estrus (heat).

ACTIVE DRUG INGREDIENT

Melengestrol acetate.....(0.0000276-0.00022%)*
Each pound contains (0.125-1.0) mg melengestrol acetate (select one)*

GUARANTEED ANALYSIS

Crude Protein (Min).....₁ _____ %
NPN (Max)..... _____ %
Crude Fat (Min)..... _____ %
Crude Fiber (Max)..... _____ %
Calcium (Min)..... _____ %
Calcium (Max)..... _____ %
Phosphorus (Min)..... _____ %
Salt (Min).....₂ _____ %
Salt (Max).....₂ _____ %
Sodium (Min).....₃ _____ %
Sodium (Max).....₃ _____ %
Potassium (Min)....._{2,4} _____ %
Vitamin A (Min)....._{2,4} _____ I.U./lb

¹ When added.

² If added.

³ Shall be guaranteed only when total sodium exceeds that furnished by the maximum salt guarantee.

⁴ Other than precursors of Vitamin A.

INGREDIENTS

Each ingredient must be named in accordance with the names and definitions adopted by the Association of American Feed Control Officials.

*Final printed label of Type C Medicated Feed must bear a single drug concentration of this drug.

FEEDING DIRECTIONS

Must be top dressed or mixed with a Type C feed containing 6.8 to 24 g/ton zilpaterol (to provide 60 mg/head/day via component feed).

Feed at the rate of 0.5 to 2.0 lb/head/day (specify one level) to provide 0.25 to 0.5 mg melengestrol acetate/head/day (specify one level). Feed melengestrol acetate in this combination for the final 20 to 40 days on feed.

MIXING DIRECTIONS

Mixing directions for liquid Type C feeds stored in recirculation tank systems are: "Recirculate immediately prior to use for no less than 10 minutes, moving not less than 1 percent of the tank contents from the bottom of the tank to the top. Recirculate daily, as directed in this paragraph even when the Type C feed is not used." Mixing directions for liquid Type C feeds stored in mechanical, air or other agitation-type tank systems are: "Agitate immediately prior to use for not less than 10 minutes, creating a turbulence at the bottom of the tank that is visible at the top. Agitate daily, as directed in this paragraph, even when the Type C feed is not used."

CAUTION: Melengestrol acetate is not effective in steers and spayed heifers. Withdrawal periods of three to five days or more should be avoided to prevent the possibility that the heifers may come into estrus (heat) at loading time.

MANUFACTURED BY
BLUE BIRD FEED MILL
Any town, USA 12345

Net Weight lb (kg) on bag or bulk

Batch (Lot or Control) No.:

(NADA 141-284) Nov 24 2014