

**BLUEBIRD FEED COMPANY
BLUEBIRD SWINE FEED
Avilamycin Type C Medicated Feed**

FOR USE IN SWINE ONLY

CAUTION: Federal law limits this drug to use under the professional supervision of a licensed veterinarian. Animal feed bearing or containing this veterinary feed directive drug shall be fed to animals only by or upon a lawful veterinary feed directive issued by a licensed veterinarian in the course of the veterinarian's professional practice.

INDICATION

For the reduction in incidence and overall severity of diarrhea in the presence of pathogenic *Escherischia coli* in groups of weaned pigs.

ACTIVE DRUG INGREDIENT

Avilamycin..... 73 grams per ton.

GUARANTEED ANALYSIS

Crude Protein, min.....	_____	%
Lysine, min.....	_____	%
Crude Fat, min.....	_____	%
Crude Fiber, max.....	_____	%
Calcium, min.....	_____	%
Calcium, max.....	_____	%
Phosphorus, min.....	_____	%
Salt ^a , min.....	_____	%
Salt ^a , max.....	_____	%
Sodium ^b , min.....	_____	%
Sodium ^b , max.....	_____	%
Selenium, min.....	_____	ppm
Zinc, min.....	_____	ppm

^a Guarantee required only when nutrient added except when the feed is intended, represented or serves as a principal source of the nutrient.

^b Sodium guarantee required only when total sodium exceeds that furnished by the maximum salt guarantee.

INGREDIENTS

Each ingredient must be named in accordance with the names and definitions adopted by the Association of American Feed Control Officials (AAFCO).

FEEDING DIRECTIONS

Feed at 73 grams per ton of Type C medicated feed (80 ppm) as the sole ration for 21 to 42* consecutive days to pigs that are at risk of developing, but not yet showing clinical signs of, diarrhea in the presence of pathogenic *Escherichia coli*.

*Final printed label on formulated Type C medicated feed must bear a single number of days, as provided on the Veterinary Feed Directive.

Caution:

To assure responsible antimicrobial drug use in pigs, do not administer to pigs 14 weeks of age or older. Do not administer medicated feed containing avilamycin to pigs for more than a lifetime total of 42 days.

Avilamycin has not been demonstrated to be effective in pigs showing clinical signs of diarrhea prior to the start of medication.

The safety of avilamycin has not been established in swine intended for breeding purposes.

Veterinary Feed Directive (VFD) expiration date must not exceed 90 days from the date of issuance. VFDs for avilamycin shall not be refilled.

WARNINGS



Residue Warning: No withdrawal period is required.



Lot Number: _____

Manufactured On: _____

MANUFACTURED BY
BLUE BIRD FEED MILL
Any town, USA 12345

Net Weight lb (kg) on bag or bulk