

**AUREO/BOV COCCI — CATTLE FEED**  
**TYPE C Medicated Feed**  
(chlortetracycline and lasalocid Type C medicated feed)

**Caution:** Federal law restricts medicated feed containing this veterinary feed directive (VFD) drug to use by or on the order of a licensed veterinarian.

For the control of coccidiosis caused by *Eimeria bovis* and *Eimeria zuernii* and for the treatment of bacterial enteritis caused by *E. coli* and bacterial pneumonia caused by *P. multocida* organisms susceptible to chlortetracycline in cattle weighing up to 800 pounds.

**ACTIVE DRUG INGREDIENTS\***

Chlortetracycline <sup>a</sup> .....	.....500 to 4000 g/ton**
Lasalocid <sup>b</sup> (as lasalocid sodium).....	.....30 to 181.8 g/ton**

**GUARANTEED ANALYSIS**

Crude protein, not less than .....	%
(This includes not more than _____% equivalent crude protein from non-protein nitrogen) <sup>1</sup>	
Crude fat, not less than.....	%
Crude fiber, not more than.....	%
Calcium, not less than .....	%
Calcium, not more than.....	%
Phosphorus, not less than.....	%
Potassium, not less than.....	%
Salt <sup>1</sup> , not less than .....	%
Salt <sup>1</sup> , not more than.....	%
Sodium <sup>2</sup> , not less than .....	%
Sodium <sup>2</sup> , not more than .....	%
Vitamin A <sup>1</sup> , not less than .....	IU/lb

<sup>1</sup>If added

<sup>2</sup>Shall be guaranteed only when total sodium exceeds that furnished by the maximum salt guarantee.

**INGREDIENTS**

Each ingredient must be specifically named in accordance with the names and definitions approved by AAFCO.

**FEEDING DIRECTIONS**

Hand feed continuously for not more than 5 days to provide 10 mg chlortetracycline per lb. body weight per day and 1 mg lasalocid per 2.2 lb. body weight per day to cattle weighing up to 800 lb. with a maximum of 360 mg of lasalocid per head per day. The following table provides examples of feeding rates and amounts:

Body Wt. Lb.	Feed Consumption		Chlortetracycline		Lasalocid	
	% Body Wt.	lb/head/day	Amount mg/lb BW/day	Feeding Rate g/ton	Amount mg/head/day	Feeding Rate g/ton
200	0.5	1	10	4000	90.9	181.8
	1	2	10	2000	90.9	90.9
	2	4	10	1000	90.9	45.5
	3	6	10	667	90.9	30.3
800	0.5	4	10	4000	360	180
	1	8	10	2000	360	90
	2	16	10	1000	360	45
	3	24	10	667	360	30

**Caution**

Safety of lasalocid in unapproved species has not been established. Do not allow horses or other equines access to feeds containing lasalocid.

► **Warning:** No withdrawal period is required. A withdrawal period has not been established for this product in pre-ruminating calves. Do not use in calves to be processed for veal.

Lot Number (if applicable): \_\_\_\_\_

**MANUFACTURED BY**

Blue Bird Feed Company  
Robin, Indiana 11111

**NET WEIGHT ON BAG OR BULK**

\*Final printed label on formulated Type C medicated feed must bear a single concentration for each drug.  
\*\*The ratio between chlortetracycline and lasalocid must be 22:1.

<sup>a</sup>Aureomycin® is the proprietary name of chlortetracycline Type A medicated article (NADA 048-761).

<sup>b</sup>BOVATEC is the proprietary name of lasalocid sodium Type A medicated article (NADA 096-298).

Zoetis Inc. [22 August 2017]