

**DAIRY AND BEEF CATTLE FEED
FBZ
TYPE B MEDICATED FEED**

INDICATIONS

For the removal and control of:

Lungworms: (*Dictyocaulus viviparus*)

Stomach worms: Barberpole worms (*Haemonchus contortus*), brown stomach worms (*Ostertagia ostertagi*), small stomach worms (*Trichostrongylus axei*)

Intestinal worms: Hookworms (*Bunostomum phlebotomum*), thread-necked intestinal worms (*Nematodirus helvetianus*), small intestinal worms (*Cooperia punctata* & *C. oncophora*)

Bankrupt worms: (*Trichostrongylus colubriformis*)

Nodular worms: (*Oesophaogostomum radiatum*)

ACTIVE DRUG INGREDIENT

Fenbendazole	1,000 – 17,740 g/ton
--------------	----------------------

GUARANTEED ANALYSIS

Crude Protein,	not less than	_____	%	
Crude Fat,	not less than	_____	%	
Crude Fiber,	not less than	_____	%	

INGREDIENTS

Each ingredient must be named in accordance with the names and definitions adopted by the Association of American Feed Control Officials Publication (AAFCO Official Publication).

MIXING DIRECTIONS

Mix 22.55 to 112.74 pounds with 1977.45 to 1887.26 pounds of feed ingredients to manufacture Type C medicated feeds containing 200 to 1000 grams of fenbendazole per ton. The resulting Type C medicated feed is to be fed at the rate of 0.454 to 2.27 pounds per 100 pounds of body weight to provided 2.27 milligrams fenbendazole per pound body weight.

WARNING: Cattle must not be slaughtered within 13 days following last treatment. For dairy cattle, the milk discard time is zero hours. A withdrawal period has not been established for this product in pre-ruminating calves. Do not use in calves to be processed for veal.

**Manufactured by
Blue Bird Feed Mills
Any town, Indiana, 12345
NET WEIGHT ON BAG OR INVOICE
NADA # 131-675, Approved by FDA**

Lot No.

Oct 2014