

**DAIRY AND BEEF CATTLE FEED  
FBZ  
TYPE C MEDICATED FEED**

**INDICATIONS**

For the removal and control of:

**Lungworms:** (*Dictyocaulus viviparus*)

**Stomach worms:** Barberpole worms (*Haemonchus contortus*), brown stomach worms (*Ostertagia ostertagi*), small stomach worms (*Trichostrongylus axei*)

**Intestinal worms:** Hookworms (*Bunostomum phlebotomum*), thread-necked intestinal worms (*Nematodirus helvetianus*), small intestinal worms (*Cooperia punctata* & *C. oncophora*)

**Bankrupt worms:** (*Trichostrongylus colubriformis*)

**Nodular worms:** (*Oesophaogostomum radiatum*)

**ACTIVE DRUG INGREDIENT**

Fenbendazole	200 – 1,000 g/ton
--------------	-------------------

**GUARANTEED ANALYSIS**

Crude Protein,	not less than	_____ %
Crude Fat,	not less than	_____ %
Crude Fiber,	not less than	_____ %

**INGREDIENTS**

Each ingredient must be named in accordance with the names and definitions adopted by the Association of American Feed Control Officials Publication (AAFCO Official Publication).

**FEEDING DIRECTIONS**

Feed at the rate of 0.454 to 2.27 pounds per 100 pounds of body weight to provided 2.27 milligrams fenbendazole per pound body weight.

**WARNING:** Cattle must not be slaughtered within 13 days following last treatment. For dairy cattle, the milk discard time is zero hours. A withdrawal period has not been established for this product in pre-ruminating calves. Do not use in calves to be processed for veal.

**Manufactured by  
Blue Bird Feed Mills  
Any town, Indiana, 12345  
NET WEIGHT ON BAG OR INVOICE**

NADA # 131-675, Approved by FDA

Lot No.

Oct 2014