

Monensin Medicated Growing Cattle Feed

Type C Medicated Feed

For Growing Cattle on Pasture or in Dry Lot

For increased rate of weight gain; For prevention and control of coccidiosis caused by *Eimeria bovis* and *Eimeria zuernii* in growing cattle on pasture or in dry lot (stocker and feeder cattle and dairy and beef replacement heifers).

Active Drug Ingredient

Monensin, USP15 to 400 g/ton*

Guaranteed Analysis

Crude Protein, not less than ¹	_____	%
Non-Protein Nitrogen (NPN) ¹ , not more than.....	_____	%
Crude Fat, not less than.....	_____	%
Crude Fiber, not more than.....	_____	%
ADF, max %.....	_____	%
Calcium, not less than.....	_____	%
Calcium, not more than.....	_____	%
Phosphorus, not less than.....	_____	%
Salt ² , not less than.....	_____	%
Salt ² , not more than.....	_____	%
Sodium ³ , not less than.....	_____	%
Sodium ³ , not more than.....	_____	%
Potassium, not less than.....	_____	%
Selenium ⁴ , min ppm.....	_____	%
Vitamin A ^{2,5} , not less than.....	_____	%

¹ When added

² If added

³ Shall be guaranteed only when total sodium exceeds that furnished by the maximum salt guarantee

⁴ Shall be stated in ppm when less than 10,000 ppm and in percentage when concentration is 10,000 ppm (1%) or greater

⁵ Other than precursors of Vitamin A

Ingredients

Each ingredient must be named in accordance with the names and definitions adopted by the Association of American Feed Control Officials.

* Final printed label on formulated Type C medicated feed must bear a single drug concentration.

Feeding Directions

Feed the appropriate amount of Type C Medicated Feed to cattle to provide 50 to 200 mg monensin/hd/day. During the first 5 days, cattle should receive no more than 100 mg monensin/day contained in not less than 1 lb. of feed.

For increased rate of weight gain: Feed at a rate of not less than 50 nor more than 200 mg monensin/hd/day in not less than 1 lb. of medicated feed;

-OR-

After the 5th day, feed at the rate of 400 mg monensin/hd/day every other day in not less than 2 lbs. of feed.

For coccidiosis prevention and control: Feed at the rate of 0.14 to 0.42 mg monensin/lb/body weight per day, depending upon severity of challenge, up to 200 mg monensin/hd/day.

Desired Intake of Monensin mg/hd/day	Concentration of Monensin In Type C Medicated Feed g/ton (mg/lb)	Recommended Feeding Level (lbs/hd/day)
60	15 (7.5)	8
60	30 (15)	4
60	60 (30)	2
100	100 (50)	2
200	200 (100)	2
150	300 (150)	1
200	400 (200)	1

Caution

Do not allow horses or other equines access to feed containing monensin. Ingestion of monensin by horses has been fatal. Monensin medicated cattle and goat feeds are safe for use in cattle and goats only. Consumption by unapproved species may result in toxic reactions. Do not exceed the levels of monensin recommended in the feeding directions, as reduced average daily gains may result. If feed refusals containing monensin are fed to other groups of cattle, the concentration of monensin in the refusals and amount of refusals fed should be taken into consideration to prevent monensin overdosing.

Warning

A withdrawal period has not been established for this product in preruminating calves. Do not use in calves to be processed for veal.

Manufactured By

Blue Bird Feed Mill Anytown, USA 12345

Net Weight lb on bag or bulk