

**Zilpaterol Finishing Cattle Feed**  
**Type C Medicated Feed**  
(zilpaterol hydrochloride Type C medicated feed)

For increased rate of weight gain, improved feed efficiency, increased carcass leanness in cattle fed in confinement for slaughter during the last 20 to 40 days on feed.

**ACTIVE DRUG INGREDIENTS**

Zilpaterol hydrochloride .....6.8 g/ton

**GUARANTEED ANALYSIS**

Crude Protein (Min) ..... \_\_\_\_\_ %  
<sup>1</sup> NPN (Max) ..... \_\_\_\_\_ %  
Crude Fat (Min) ..... \_\_\_\_\_ %  
Crude Fiber (Max) ..... \_\_\_\_\_ %  
Calcium (Min) ..... \_\_\_\_\_ %  
Calcium (Max) ..... \_\_\_\_\_ %  
Phosphorus (Min) ..... \_\_\_\_\_ %  
<sup>2</sup> Salt (Min) ..... \_\_\_\_\_ %  
<sup>2</sup> Salt (Max) ..... \_\_\_\_\_ %  
<sup>3</sup> Sodium (Min) ..... \_\_\_\_\_ %  
<sup>3</sup> Sodium (Max) ..... \_\_\_\_\_ %  
Potassium (Min) ..... \_\_\_\_\_ %  
<sup>2,4</sup> Vitamin A (Min) ..... \_\_\_\_\_ I.U./lb

<sup>1</sup> When added.

<sup>2</sup> If added.

<sup>3</sup> Shall be guaranteed only when total sodium exceeds that furnished by the maximum salt guarantee.

<sup>4</sup> Other than precursors of Vitamin A.

**INGREDIENTS**

Each ingredient must be named in accordance with the names and definitions adopted by the Association of American Feed Control Officials.

## FEEDING DIRECTIONS

Feed continuously as sole ration to cattle during the last 20 to 40 days on feed, Zilpaterol Finishing Cattle Feed Type C Medicated Feed containing 6.8 g/ton zilpaterol hydrochloride to provide 60 to 90 mg/head/day zilpaterol hydrochloride.

**WARNING:** The active ingredient in Zilmax<sup>®</sup> is zilpaterol hydrochloride, a beta<sub>2</sub>-adrenergic agonist. Not for use in humans. An anti-dust process has been applied to the drug product, Zilmax<sup>®</sup>, in order to greatly reduce inhalation risk. Extended handling tasks with the potential for dust generation require respiratory protection. Wear appropriate skin protection (e.g., impervious gloves, apron, overalls), if there is a potential for extended skin contact. Wear protective eyewear, if there is a potential for eye contact. If accidental eye contact occurs, immediately rinse with water and consult a physician.

### ► WITHDRAWAL PERIOD: 3 days ◀

**CAUTION:** Zilpaterol hydrochloride is not for use in animals intended for breeding. Do not allow horses or other equines access to feed containing zilpaterol. Do not use in veal calves. Not to be fed to cattle in excess of 90 mg zilpaterol/head/day in complete feed. If pen consumption of complete feed exceeds 26.5 lb/head/day (90% dry matter basis), zilpaterol should not be fed in complete feed.

**YOU MAY NOTICE:** Animals receiving zilpaterol hydrochloride may exhibit increased respiratory rate as well as elevated levels of creatine phosphokinase (CPK) and creatinine.

**MANUFACTURED BY**  
BLUE BIRD FEED MILL  
Any town, USA 12345

**Net Weight lb (kg) on bag or bulk**

**Batch (Lot or Control) No.:**

(NADA 141-284) Apr 16, 2015