

Zilpaterol Finishing Cattle Feed
Type C Medicated Component Feed
(zilpaterol hydrochloride Type C medicated feed)

For increased rate of weight gain, improved feed efficiency, and increased carcass leanness in cattle fed in confinement for slaughter during the last 20 to 40 days on feed.

ACTIVE DRUG INGREDIENT

Zilpaterol hydrochloride 6.8 – 24 g/ton*

GUARANTEED ANALYSIS

Crude Protein (Min).....	_____ %
NPN ¹ (Max).....	_____ %
Crude Fat (Min).....	_____ %
Crude Fiber (Max).....	_____ %
Calcium (Min).....	_____ %
Calcium (Max).....	_____ %
Phosphorus (Min).....	_____ %
Salt ² (Min).....	_____ %
Salt ² (Max).....	_____ %
Sodium ³ (Min).....	_____ %
Sodium ³ (Max).....	_____ %
Potassium (Min).....	_____ %
Vitamin A ^{2,4} (Min).....	_____ I.U./lb

¹ When added.

² If added.

³ Shall be guaranteed only when total sodium exceeds that furnished by the maximum salt guarantee.

⁴ Other than precursors of Vitamin A.

INGREDIENTS

Each ingredient must be named in accordance with the names and definitions adopted by the Association of American Feed Control Officials.

*Final printed label on formulated Type C Medicated Feed must bear a single drug concentration of this drug.

FEEDING DIRECTIONS

Feed continuously to cattle during the last 20 to 40 days on feed, Zilpaterol Finishing Cattle Feed Type C Medicated Component Feed containing 6.8 to 24 g/ton zilpaterol hydrochloride to provide 60 mg/head/day zilpaterol hydrochloride.

The following table gives examples of how some Type C medicated component feed concentrations can be fed at different target consumption levels to deliver zilpaterol hydrochloride at 60 mg/head/day.

Zilpaterol hydrochloride in Type C Medicated Component Feed ¹	Type C Medicated Component Feed Consumption	Zilpaterol hydrochloride Consumed
g/ton [*]	lb/hd/day	mg/hd/day
24	5	60
10	12	60

¹ Based on 90% dry matter basis.

^{*} The maximum amount of Type C medicated component feed in a daily feeding must not exceed 17.6 lb, when zilpaterol hydrochloride is provided in the minimum approved concentration of 6.8 g/ton.

WARNING: The active ingredient in Zilmax[®] is zilpaterol hydrochloride, a beta₂-adrenergic agonist. Not for use in humans. An anti-dust process has been applied to the drug product, Zilmax[®], in order to greatly reduce inhalation risk. Extended handling tasks with the potential for dust generation require respiratory protection. Wear appropriate skin protection (e.g., impervious gloves, apron, overalls), if there is a potential for extended skin contact. Wear protective eyewear, if there is a potential for eye contact. If accidental eye contact occurs, immediately rinse with water and consult a physician.

► **WITHDRAWAL PERIOD: 3 days** ◀

CAUTION: Zilpaterol hydrochloride is not for use in animals intended for breeding. Do not allow horses or other equines access to feed containing zilpaterol. Do not use in veal calves.

YOU MAY NOTICE: Animals receiving zilpaterol hydrochloride may exhibit increased respiratory rate as well as elevated levels of creatine phosphokinase (CPK) and creatinine.

MANUFACTURED BY
BLUE BIRD FEED MILL
Any town, USA 12345

Net Weight lb (kg) on bag or bulk

Batch (Lot or Control) No.:

(NADA 141-284) Apr 16, 2015