BLUEBIRD FEED COMPANY BLUE BIRD SWINE FEED

Type B Medicated Feed for Swine

Caution: Federal law restricts medicated feed containing this veterinary feed directive (VFD) drug to use by or on the order of a licensed veterinarian.

Indication

For the control of swine respiratory disease (SRD) associated with *Actinobacillus pleuropneumoniae, Pasteurella multocida, Streptococcus suis,* and *Bordetella bronchiseptica* in groups of swine in buildings experiencing an outbreak of SRD.

Active Drug Ingredient

Guaranteed Analysis

Crude Protein (Min) %	Calcium (min and max)%
Crude Fat (Min)%	Phosphorous (min)%
Crude Fiber (Max)%	Salt (min and max)% ¹
Total Ash (Max)%	Selenium (min)ppm
Moisture (Max) %	Zinc (min)ppm
Lysine (min)%	¹ If Added

Ingredients

Each ingredient must be named in accordance with the names and definitions adopted by the Association of American Feed Control Officials (AAFCO).

Mixing Instructions

Thoroughly mix 20 lb of this Type B medicated Feed for Swine with 1,980 lb of feed to provide Type C Medicated Feed containing 182 g florfenicol per ton, as shown below:

Starting concentration of florfenicol	Amount of florfenicol Type B	Resulti	ng concentration	in
Type B Medicated Feed for Swine	Medicated Feed to add per ton	Type	C Medicated Feed	l
grams/pound	pounds	grams/ton	grams/pound	ppm
9.1	20	182	0.091	200

Feeding Directions

Feed the resulting Type C medicated feed as the sole ration for 5 consecutive days to swine to deliver 10 mg florfenicol per kg body weight per day.

Caution: Feed containing florfenicol shall not be fed to pigs for more than 5 days. Following administration, pigs should be reevaluated by a licensed veterinarian before reinitiating a further course of therapy. The expiration date for VFD for Nuflor® (florfenicol) must not exceed 90 days from the date of issuance. VFD for Nuflor® (florfenicol) shall not be refilled. The effects of florfenicol on swine reproductive performance, pregnancy, and lactation have not been determined.

Warning: Avoid inhalation, oral exposure, and direct contact with skin or eyes. Operators mixing and handling Nuflor® Type A medicated article for Swine and Type B medicated feed for Swine should use protective clothing, gloves, goggles and a NIOSH-approved dust mask. Wash thoroughly with soap and water after handling. If accidental eye contact occurs, immediately rinse thoroughly with water. If irritation persists, seek medical attention. Not for human consumption. Keep out of reach of children. The Safety Data Sheet (SDS) contains more detailed occupational safety information. To obtain more information or to report adverse effects call 1-800-211-3573. For customer service or a copy of the SDS, call 1-800-521-5767.

Storage Conditions: Store in a cool, dry place. Avoid excessive moisture and heat.



Residue Warning: Feeds containing florfenicol must be withdrawn 13 days prior to slaughter.



Manufactured By: Bluebird Feed M	ill, Robin, IN 46813
NET WEIGHT 20 lbs (9.1 kg)	Lot No.:



Nuflor $^{\mathbb{R}}$ is a registered trademark of Intervet Inc, (d/b/a) Merck Animal Health.

Copyright©2015. Intervet Inc., Madison, NJ 07940, USA. Manufactured for Intervet Inc. All rights reserved. 09/2015

