

Monensin Medicated Turkey Feed
(monensin Type C medicated feed)

INDICATION

For the prevention of coccidiosis in turkeys caused by *Eimeria adenoeides*, *E.meleagrimitis* and *E. gallopavonis*.

ACTIVE DRUG INGREDIENT

Monensin^a, USP54 - 90 g/ton*

GUARANTEED ANALYSIS

Crude Protein, not less than.....	_____	%
Lysine, not less than.....	_____	%
Methionine, not less than.....	_____	%
Crude Fat, not less than.....	_____	%
Crude Fiber, not more than.....	_____	%
Calcium, not less than.....	_____	%
Calcium, not more than.....	_____	%
Phosphorus, not less than.....	_____	%
Salt ¹ , not less than.....	_____	%
Salt ¹ , not more than.....	_____	%
Sodium ² , not less than.....	_____	%
Sodium ² , not more than.....	_____	%

¹If added.

²Shall be guaranteed only when total sodium exceeds that furnished by the maximum salt guarantee.

INGREDIENTS

Each ingredient must be named in accordance with the names and definitions adopted by the Association of American Feed Control Officials.

FEEDING DIRECTIONS

Feed 54 - 90 g/ton of complete feed. For growing turkeys only, feed continuously as the sole ration. The optimum level depends upon the severity of coccidiosis exposure.

WARNING

Do not feed to laying chickens. Do not feed to chickens over 16 weeks of age.

CAUTION

Do not allow horses, other equines, mature turkeys, or guinea fowl access to feed containing monensin. Ingestion of monensin by horses and guinea fowl has been fatal. Some strains of turkey coccidia may be monensin tolerant or resistant. Monensin may interfere with development of immunity to turkey coccidiosis. Not for replacement chickens intended to become broiler breeding chickens.

MANUFACTURED BY
BLUE BIRD FEED MILL
Any town, USA 12345

Net Weight lb (kg) on bag or bulk

*The medicated feed label must state a single drug concentration.

^aSourced from CobanTM, NADA# 38-878 and 130-736

Elanco, Coban and the diagonal bar are trademarks owned or licensed by Eli Lilly and Company, its subsidiaries or affiliates.