

BLUE BIRD
DCX Liquid Sheep & Goat Feed
(decoquinatate Type B liquid medicated feed)

Sheep: For the prevention of coccidiosis in young sheep caused by *Eimeria ovinoidalis*, *E. parva*, *E. bakuensis*, and *E. crandallis*. **Goats:** For the prevention of coccidiosis in young goats caused by *Eimeria christensenii* and *E. ninakohlyakimovae*.

Active Drug Ingredient

Decoquinatate.....90.9 - 5440 g/ton*

Guaranteed Analysis

Crude Protein (Min).....	_____	%
NPN ^a (Max).....	_____	%
Crude Fat (Min).....	_____	%
Crude Fiber (Max).....	_____	%
Calcium ^a (Min).....	_____	%
Calcium ^a (Max).....	_____	%
Phosphorus ^a (Min).....	_____	%
Salt ^a (Min).....	_____	%
Salt ^a (Max).....	_____	%
Sodium ^b (Min).....	_____	%
Sodium ^b (Max).....	_____	%
Copper ^{c, d} (Min).....	_____	ppm or %
Copper ^{c, d} (Max).....	_____	ppm or %
Selenium ^{a, d} (Min).....	_____	ppm or %
Vitamin A ^a (Min).....	_____	IU/lb

This liquid Type B medicated feed must have a pH between 5.0 – 6.5 and contain a suspending agent to maintain viscosity of not less than 500 centipoises.

- ^a Guarantee required only when nutrient added except when the feed is intended, represented or serves as a principal source of the nutrient.
- ^b Sodium guarantee required only when total sodium exceeds that furnished by the maximum salt guarantee.
- ^c Copper minimum and maximum guarantee for sheep required when added or level exceeds 20 ppm.
- ^d Guarantee shall be stated in ppm when less than 10,000 ppm and in percentage when concentration is 10,000 ppm (1%) or greater.

Ingredients

Ingredients as defined by AAFCO.

Mixing Directions

Mix this liquid Type B medicated feed with non-medicated feed ingredients to manufacture one ton of Type C medicated feed containing 12.9 to 90.8 grams decoquinatate.

The following table provides examples of *mixing* rates:

Type B Decoquinatate concentration (g/ton)	Type B per ton of Type C (lb)	Non-medicated feed per ton of Type C (lb)	Type C Decoquinatate concentration (g/ton)
568	80	1920	22.7
2270	40	1960	45.4
5440	4.75	1995.25	12.9
5440	33.4	1996.6	90.8

The resulting Type C medicated feed should be fed for at least 28 days during periods of exposure to coccidiosis or when it is likely to be a hazard. The concentration of decoquinatate in the final Type C feed will vary with body weight and feed consumption to provide 22.7 mg per 100 pounds of body weight (0.5 mg/kg) per day.

Warning: Do not feed to sheep and goats producing milk for human consumption.

Caution: Do not use in feeds containing bentonite.

Manufactured by
Blue Bird Feed Mill
City, State, Zip

NET WEIGHT ON BAG OR BULK

Expiration Date (if applicable) _____

Lot Number (if applicable)) _____

*The final printed feed label should list only a single drug concentration.
039417_12212015
Zoetis Inc.