

**BLUE BIRD**  
**DCX Top-dress Sheep & Goat Feed**  
(decoquinate Type B medicated feed)

**Sheep:** For the prevention of coccidiosis in young sheep caused by *Eimeria ovinoidalis*, *E. parva*, *E. bakuensis*, and *E. crandallis*. **Goats:** For the prevention of coccidiosis in young goats caused by *Eimeria christensenii* and *E. ninakohlyakimovae*.

**Active Drug Ingredient**

Decoquinate.....535.8 - 5440 g/ton\*

**Guaranteed Analysis**

Crude Protein (Min).....	_____	%
NPN <sup>a</sup> (Max).....	_____	%
Crude Fat (Min).....	_____	%
Crude Fiber (Max).....	_____	%
Calcium <sup>a</sup> (Min).....	_____	%
Calcium <sup>a</sup> (Max).....	_____	%
Phosphorus <sup>a</sup> (Min).....	_____	%
Salt <sup>a</sup> (Min).....	_____	%
Salt <sup>a</sup> (Max).....	_____	%
Sodium <sup>b</sup> (Min).....	_____	%
Sodium <sup>b</sup> (Max).....	_____	%
Copper <sup>c, d</sup> (Min).....	_____	ppm or %
Copper <sup>c, d</sup> (Max).....	_____	ppm or %
Selenium <sup>a, d</sup> (Min).....	_____	ppm or %
Vitamin A <sup>a</sup> (Min).....	_____	IU/lb

<sup>a</sup> Guarantee required only when nutrient added except when the feed is intended, represented or serves as a principal source of the nutrient.

<sup>b</sup> Sodium guarantee required only when total sodium exceeds that furnished by the maximum salt guarantee.

<sup>c</sup> Copper minimum and maximum guarantee for sheep required when added or level exceeds 20 ppm.

<sup>d</sup> Guarantee shall be stated in ppm when less than 10,000 ppm and in percentage when concentration is 10,000 ppm (1%) or greater.

**Ingredients**

Ingredients as defined by AAFCO.

**Mixing Directions**

Mix this Type B medicated feed with non-medicated feed ingredients to manufacture one ton of a Type C supplement containing 90.9 to 535.7 grams of decoquinate.

The following table provides examples of mixing rates:

Type B Decoquinate concentration (g/ton)	Type B per ton of Type C (lb)	Non-medicated feed per ton of Type C (lb)	Type C Decoquinate concentration (g/ton)
2270	80	1920	90.9
5440	50	1950	136.0
5440	167	1833	454.2

The resulting Type C medicated feed should be top-dressed on the daily ration for at least 28 days during periods of exposure to coccidiosis or when it is likely to be a hazard. The concentration of decoquinate in the final Type C feed will vary with body weight and feed consumption to provide 22.7 mg per 100 pounds of body weight (0.5 mg/kg) per day.

**Warning:** Do not feed to sheep and goats producing milk for human consumption.

**Caution:** Do not use in feeds containing bentonite.

**Manufactured by:**

Blue Bird Feed Mill  
City, State, Zip

**NET WEIGHT ON BAG OR BULK**

Expiration Date (if applicable) \_\_\_\_\_

Lot Number (if applicable) ) \_\_\_\_\_

\*The final printed feed label should list only a single drug concentration.

039417\_12212015  
Zoetis Inc.