

**BLUE BIRD**  
**DCX Sheep & Goat Feed**  
(decoquinatate Type B medicated feed)

**Sheep:** For the prevention of coccidiosis in young sheep caused by *Eimeria ovinoidalis*, *E. parva*, *E. bakuensis*, and *E. crandallis*.

**Goats:** For the prevention of coccidiosis in young goats caused by *Eimeria christensenii* and *E. ninakohlyakimovae*.

**Active Drug Ingredient**

Decoquinatate.....90.9 - 5440 g/ton\*

**Guaranteed Analysis**

|                                     |       |          |
|-------------------------------------|-------|----------|
| Crude Protein (Min).....            | _____ | %        |
| NPN <sup>a</sup> (Max).....         | _____ | %        |
| Crude Fat (Min).....                | _____ | %        |
| Crude Fiber (Max).....              | _____ | %        |
| Calcium <sup>a</sup> (Min).....     | _____ | %        |
| Calcium <sup>a</sup> (Max).....     | _____ | %        |
| Phosphorus <sup>a</sup> (Min).....  | _____ | %        |
| Salt <sup>a</sup> (Min).....        | _____ | %        |
| Salt <sup>a</sup> (Max).....        | _____ | %        |
| Sodium <sup>b</sup> (Min).....      | _____ | %        |
| Sodium <sup>b</sup> (Max).....      | _____ | %        |
| Copper <sup>c, d</sup> (Min).....   | _____ | ppm or % |
| Copper <sup>c, d</sup> (Max).....   | _____ | ppm or % |
| Selenium <sup>a, d</sup> (Min)..... | _____ | ppm or % |
| Vitamin A <sup>a</sup> (Min).....   | _____ | IU/lb    |

<sup>a</sup> Guarantee required only when nutrient added except when the feed is intended, represented or serves as a principal source of the nutrient.

<sup>b</sup> Sodium guarantee required only when total sodium exceeds that furnished by the maximum salt guarantee.

<sup>c</sup> Copper minimum and maximum guarantee for sheep required when added or level exceeds 20 ppm.

<sup>d</sup> Guarantee shall be stated in ppm when less than 10,000 ppm and in percentage when concentration is 10,000 ppm (1%) or greater.

**Ingredients**

Ingredients as defined by AAFCO.

**Mixing Directions**

Mix this Type B medicated feed with non-medicated feed ingredients to manufacture one ton of complete feed or milk replacer containing 12.9 to 90.8 grams decoquinatate.

The following table provides examples of mixing rates:

| Type B Decoquinatate concentration (g/ton) | Type B per ton of Type C (lb) | Non-medicated feed per ton of Type C (lb) | Type C Decoquinatate concentration (g/ton) |
|--|-------------------------------|---|--|
| 568  | 80                            | 1920                                      | 22.7                                       |
| 2270                                       | 40                            | 1960                                      | 45.4                                       |
| 5440                                       | 4.75                          | 1995.25                                   | 12.9                                       |
| 5440                                       | 33.4                          | 1996.6                                    | 90.8                                       |

The resulting Type C medicated feed should be fed for at least 28 days during periods of exposure to coccidiosis or when it is likely to be a hazard. The concentration of decoquinatate in the final Type C feed will vary with body weight and feed consumption to provide 22.7 mg per 100 pounds of body weight (0.5 mg/kg) per day.

**Warning:** Do not feed to sheep and goats producing milk for human consumption.

**Caution:** Do not use in feeds containing bentonite.

**Manufactured by:**

Blue Bird Feed Mill

City, State, Zip

**NET WEIGHT ON BAG OR BULK**

\*The final printed feed label should list only a single drug concentration.

039417\_12212015

Zoetis Inc.